

Integrable viscous conservation laws

Alessandro Arsie*, Paolo Lorenzoni**, Antonio Moro**

*Department of Mathematics and Statistics

University of Toledo, 2801 W. Bancroft St., 43606 Toledo, OH, USA

**Dipartimento di Matematica e Applicazioni

Università di Milano-Bicocca, Via Roberto Cozzi 53, I-20125 Milano, Italy

Abstract

We propose an extension of Dubrovin's perturbative approach to the study of normal forms for non-Hamiltonian integrable scalar conservation laws. The explicit computation of the first few corrections leads to the conjecture that such normal forms are parametrized by one single functional parameter, named viscous central invariant. A constant valued viscous central invariant corresponds to the well-known Burgers hierarchy. The case of a linear viscous central invariant provides a viscous analog of the Camassa-Holm equation, that formerly appeared as a reduction of a two-component generalization of the Camassa-Holm equation. We write explicitly the negative and positive hierarchy associated with this equation and prove the integrability showing that they can be mapped respectively into the heat hierarchy and its negative counterpart, named Klein-Gordon hierarchy. A local well-posedness theorem for periodic initial data is also proven. Finally, we explicitly construct asymptotic solutions obtained by quasi-Miura transformations and discuss Dubrovin's Universality conjecture for the local behaviour of the solutions near the point of gradient catastrophe.

1 Introduction

In [12] B. Dubrovin proposed a perturbative approach to the study of integrable Hamiltonian evolutionary equations (or systems of equations) of the form

$$u_t = A(u)u_x + \epsilon[B_1(u)u_{xx} + B_2(u)u_x^2] + \epsilon^2[C_1(u)u_{xxx} + C_2(u)u_x u_{xx} + C_3(u)u_x^3] + \mathcal{O}(\epsilon^3). \quad (1.1)$$

where ϵ is a small formal expansion parameter. If the unperturbed equation obtained for $\epsilon = 0$ is integrable and Hamiltonian (this is always true in the scalar case), Dubrovin's method provides an effective tool to construct perturbative dispersive equations of the form (1.1) preserving integrability and Hamiltonian properties. The r.h.s. of (1.1) is a formal expansion, it contains in principle infinitely many terms and no additional assumptions concerning convergence are usually enforced. It is natural to classify equations of the form (1.1) up to the group of Miura transformations defined via formal series of the form

$$u \rightarrow v = f(u) + \epsilon[g(u)u_x] + \epsilon^2[h_1(u)u_{xx} + h_2(u)u_x^2] + \dots \quad (1.2)$$

Since most of physical applications involving equations of the form (1.1) contain a finite number of terms, one would be tempted to work with truncated Miura transformations only. This choice turns out to be ineffective and the reason is two-fold. First of all, an infinite formal expansion is needed, in general, to define the inverse of a Miura transformation, truncated or not: if one does not allow infinite formal expansion, then the set of truncated Miura transformations do not form a group. Secondly, the restriction to finite expansions will exclude from the analysis all those non-evolutionary equations that can be put in the evolutionary form (1.1). Let us consider for example the celebrated Camassa-Holm equation

$$u_t - \epsilon^2 u_{xxt} = -3uu_x + \epsilon^2(uu_{xxx} + 2u_x u_{xx}). \quad (1.3)$$

Introducing the transformation

$$v = u - \epsilon^2 u_{xx}$$

its formal inversion is given by an infinite formal series that allows to re-write the equation (1.3) in the evolutionary form (1.1) with respect to the variable v .

However in a number of cases truncated equations might have their own interest and in particular those equations possessing infinitely many approximate symmetries [24, 33].

In this paper we will extend Dubrovin's approach to the case of integrable conservation laws of the form

$$u_t = \partial_x \{u^2 + \epsilon[a(u)u_x] + \epsilon^2[b_1(u)u_{xx} + b_2(u)u_x^2] + \mathcal{O}(\epsilon^3)\}. \quad (1.4)$$

Equations of this form are not necessarily Hamiltonian. For instance, in the scalar case, the operator ∂_x defines a Poisson structure on the space of functionals $F[u] = \int f(u, u_x, u_{xx}, \dots) dx$, but in general the current

$$u^2 + \epsilon[a(u)u_x] + \epsilon^2[b_1(u)u_{xx} + b_2(u)u_x^2] + \mathcal{O}(\epsilon^3)$$

is not the variational derivative of a functional $F[u]$. To overcome this difficulty we consider the extension of the Hamiltonian formalism to 1-forms introduced in [1]. Within this framework, currents are viewed as 1-forms in a suitable jet space on which the operator ∂_x defines a Poisson structure. Hence, the extension of Dubrovin perturbative scheme is immediate apart from the choice of the class of admissible formal Miura transformations. There are two natural options:

1. To consider canonical transformations as in [12] only. This choice is motivated by results in [19] that guarantees any deformation of the operator ∂_x to be eliminated by a Miura transformation.
2. To consider Miura transformations that preserve the form of the equation (1.4). This class is clearly wider and plays a crucial role in the elimination of few inessential functional parameters. This is the reason why we have chosen to work with this second option.

An alternative approach to the classification of integrable hierarchies, based on the nonlinear perturbations to linear equations, has been developed for instance in [27], [28], [29], [2]. This is the so called symmetry approach.

The main results of this paper can be summarized as follows:

- We classify up to the fifth order in the deformation parameter scalar integrable conservation laws of the form (1.4). We will focus on the viscous case corresponding to $a(u) \neq 0$ and for which the corresponding currents are not exact 1-forms. It turns out that, up to the class Miura transformations preserving the form of the equation (1.4), all deformations are uniquely determined by the function $a(u)$ named viscous central invariant. For this reason, we refer to the corresponding equations as *integrable viscous conservation laws*. Dispersive deformations obtained for $a(u) = 0$ and $b_1(u) \neq 0$ have been already studied in [12] in the case of Hamiltonian conservation laws.
- We observe that except for the case of constant viscous central invariant where the right hand side of (1.4) becomes finite and gives the well known Burgers equation, the generic integrable conservation law contains infinitely many terms. We show that the case of a linear viscous central invariant $a(u) = u$ corresponds to the equation

$$u_t - \epsilon u_{xt} = \partial_x (u^2 - \epsilon u u_x), \quad (1.5)$$

formerly derived as a non-Hamiltonian reduction of two-component generalizations of the Camassa-Holm equation [31, 16]. A few interesting properties of equation (1.5) have been already investigated, such as the existence of a typical viscous shock dynamics associated with cliffons, i.e. weak traveling wave,

solutions [16]. Nevertheless, its integrable structure has not been yet studied in detail. We observe that, remarkably, equation (1.5) is related by a Miura transformation to the equation

$$u_t = \partial_x \left(\sum_k \epsilon^k u u_{(k)} \right) \quad (1.6)$$

which is a symmetry of the equation

$$u_{\tau_{-1}} = \partial_x \left(\frac{1}{u} + \epsilon \partial_x \frac{1}{u} \right) \quad (1.7)$$

appearing in Calogero's list [4]. We show that the equation (1.7) is the first member of the full integrable “negative” hierarchy

$$u_{\tau_{-n}} = \partial_x \left(\frac{1}{u} + \epsilon \partial_x \frac{1}{u} \right)^n (1) \quad n = 1, 2, 3, \dots \quad (1.8)$$

We also construct the positive counterpart which turns out to be non-local, that is its evolutionary form is not truncated, as it happens for the equation (1.6). It should be noted that the negative hierarchy (1.8) and its positive counterparts can be obtained as a one component reduction on the negative hierarchy recursively constructed in [16]. We provide an alternative proof of the integrability by showing that the negative and positive hierarchies can be transformed respectively into the Burgers and then the heat hierarchy and its negative counterpart. Interestingly the first member of the negative counterpart is given by the Klein-Gordon equation and for this reason we refer to it as Klein-Gordon hierarchy.

- We study the well-posedness (existence, uniqueness and continuous dependence on initial data) for the periodic Cauchy problem for the equation (1.5) (with $\epsilon = 1$) locally in time, using the general techniques developed by Kato in [18], similarly to what has been done for the Camassa-Holm equation in [7]. More specifically, we prove that the periodic Cauchy problem for the equation (1.5) is locally well-posed for $m \in H^1$, where $m = u - u_x$. It is a work in progress the determination of the conditions that guarantee the global well-posedness of the periodic Cauchy problem and the blow-up conditions in finite time, in full analogy with the results obtained for the Camassa-Holm equation [7, 6]. We note that in [15] well-posedness and blow-up properties have been studied for the two component generalization of the equation (1.5) introduced in [5] and [16].
- Using **Mathematica** [34] we investigate numerically the Cauchy problem for the equation (1.5) assuming periodic boundary conditions. This has been done by

converting the equation (1.5) into a system consisting of an evolutionary PDE together with an ODE. The numerical integrator, based on the method of lines for solving the PDEs, applies a pseudo-spectral method or a spatial discretization with finite differences of the 4th order. We perform numerical simulations with very simple initial data such as translated sinusoidal waves with different wave lengths and observe that solutions behaviour is as expected: the initial datum evolves in low viscosity regime approaching the gradient catastrophe and the steep profile so created is asymptotically dissipated. We also provide numerical evidence of the existence of initial conditions for which the solution breaks in finite time similarly to what happens for the Camassa-Holm equation in [7].

- The class of perturbed equations considered in this paper coincides up to $O(\epsilon^2)$ with the generalized Burgers equations studied in [9]. We show that near the critical point the solutions of scalar Hamiltonian conservation laws should satisfy a second order ODE which is the non-Hamiltonian analogue of the Painlevé equation arising in the description of the critical behaviour of Hamiltonian conservation laws [12]. The mentioned ODE admits the Pearcey function as a particular solution. This is consistent with the results in [21] and [9] concerning the universality of the critical behaviour of Burgers' equation and its generalization.

The paper is organized as follows. In Section 2 we give the definition of integrable scalar conservation laws and we introduce their normal form in Section 3. In Section 4 we present the perturbative classification results concerning integrable viscous conservation laws and discuss in detail the case of a linear viscous central invariant in Section 5. Section 6 is devoted to the proof of the local well-posedness for the Cauchy problem and numerical simulations are presented in Section 7. In Section 8 we review some results about the quasitriviality of evolutionary PDEs due to Liu and Zhang [23]. However, instead of characterizing the quasitriviality transformations in terms of their infinitesimal generators as in [23], we provide a direct characterization of the coefficients of the transformation. In Section 9 these results are applied to the study of the critical behaviour of solutions near the critical point.

2 Commuting flows

In the present section we introduce the main notions concerning integrability and perturbation theory of conservation laws. For the sake of simplicity we specify all definitions and properties in the case of scalar conservation laws, that is the subject

matter of this paper. The interested reader will find a rigorous and more general treatment in the paper [1].

Let

$$u_t = \partial_x \alpha(u, u_x, u_{xx}, \dots) \quad (2.1)$$

be a scalar conservation law. If we assume that the current α is a differential polynomial, then, rescaling the variables as $x \rightarrow \epsilon x, t \rightarrow \epsilon t$, we can write the right hand side of any scalar conservation law as

$$u_t = \partial_x \left(\alpha_0(u) + \sum_{k=1}^N \epsilon^k \alpha_k(u, u_x, u_{xx}, \dots) \right) \quad (2.2)$$

where α_k are differential polynomials of degree k . Here the degree is assigned according to the following rule: $\deg f(u) = 0, \deg u_{(k)} = k$. Clearly if we start from a differential polynomial, N is finite. However as mentioned above we also include the case $N = \infty$. In this case the equation (2.2) will be referred to as *formal conservation law*.

Definition 2.1 *A (formal) scalar conservation law (2.2) is said to be integrable if it admits infinitely many symmetries of the form*

$$u_\tau = \partial_x \beta(u, u_x, u_{xx}, \dots). \quad (2.3)$$

A very important subcase is given by the class Hamiltonian conservation laws. In this case the current α is the variational derivative of a suitable local functional (the Hamiltonian functional)

$$H[u] = \int h(u, u_x, \dots) dx$$

where the function h is the Hamiltonian density. In this case we have

$$\alpha = \frac{\delta H}{\delta u} = \frac{\partial h}{\partial u} - \partial_x \left(\frac{\partial h}{\partial u_x} \right) + \partial_x^2 \left(\frac{\partial h}{\partial u_{xx}} \right) + \dots$$

Two functionals $F[u]$ and $G[u]$ such that $\{F, G\} = 0$ are said to be *in involution*.

It is well-known (see e.g. [10]) that the operator ∂_x defines a Poisson bracket in the space of local functionals. Given two local functionals $F[u]$ and $G[u]$, their Poisson bracket is the local functional defined by

$$\{F[u], G[u]\} = \int \frac{\delta F}{\delta u} \partial_x \left(\frac{\delta G}{\delta u} \right) dx. \quad (2.4)$$

It was proved in [19, 8] that any local Hamiltonian operator can be reduced in the scalar case to ∂_x by means of a Miura transformation. This means that any scalar

Hamiltonian equation (w.r.t a local Hamiltonian operator) can be written as a conservation law after a suitable Miura transformation. Moreover, in the Hamiltonian case also the symmetries of the equation are Hamiltonian and the corresponding Hamiltonian functionals are in involution with the Hamiltonian functional of the original equation.

The previous analysis shows that the function α in (2.2) can be thought as a 1-form. Such a 1-form is in general non-exact and the equation is not Hamiltonian with respect to the Hamiltonian operator ∂_x .

Nevertheless, extending a well-known construction in the finite dimensional setting, it is possible to define a Poisson bracket on the space of 1-forms Λ_1 . In the scalar case, this is given by:

$$\{\alpha, \beta\} := \sum_j \partial_x^{j+1} \beta \frac{\partial \alpha}{\partial u_{(j)}} - \partial_x^{j+1} \alpha \frac{\partial \beta}{\partial u_{(j)}} = 0. \quad (2.5)$$

The above bracket satisfies the following properties (see [1] for more details):

1. If $\alpha = \delta F$ and $\beta = \delta G$, then $\{\alpha, \beta\} = \delta\{F, G\}$.
2. $\{\cdot, \cdot\}$ equips the space of 1-forms Λ_1 with a Lie algebra structure;
3. the Poisson structure induces an (anti)-homomorphism of Lie algebras between $(\Lambda_1, \{\cdot, \cdot\})$ and the space of evolutionary vector fields equipped with the Lie bracket given by the Lie commutator.

From the last property it follows that two scalar conservation laws commute if and only if the associated currents are in involution. For convenience of the reader we give an alternative elementary proof of this fact.

Proposition 2.2 *Two flows of conservation laws of the form (2.2) and (2.3) commute iff α and β are in involution with respect to the bracket (2.5).*

Proof: Let us introduce the function φ such that $u = \varphi_x$. Then, integrating once w.r.t. the variable x , equations (2.2) and (2.3) give

$$\begin{aligned} \varphi_t &= \alpha(u, u_x, u_{xx}, \dots) + f(t, \tau) \\ \varphi_\tau &= \beta(u, u_x, u_{xx}, \dots) + g(t, \tau) \end{aligned}$$

where f and g are two arbitrary functions. The request that the two equations above commute for any solution φ , i.e.

$$\partial_\tau \varphi_t = \partial_t \varphi_\tau$$

implies that the two compatibility conditions

$$\alpha_\tau = \beta_t \quad f_\tau = g_t$$

must hold separately. The second condition is identically satisfied if one choses $f = \psi_t$ and $g = \psi_\tau$. Using the chain rule together with the equations (2.2) and (2.3) the first condition can be equivalently written as follows

$$\alpha_\tau - \beta_t = \sum_j \partial_x^{j+1} \beta \frac{\partial \alpha}{\partial u_{(j)}} - \partial_x^{j+1} \alpha \frac{\partial \beta}{\partial u_{(j)}} = 0.$$

The proposition is proved. ■

As a consequence of the above proposition, we have that any integrable hierarchy of conservation laws is always defined by an infinite family of 1-forms in involution. From this perspective Hamiltonian and non-Hamiltonian conservation laws can be understood within the same framework.

Example Let us consider the Burgers hierarchy defined as

$$u_{t_n} = \partial_x \omega_n = \partial_x [(u + \partial_x)^n u], \quad n = 0, 1, 2, \dots \quad (2.6)$$

The flows associated with ω_n are in involution w.r.t. the Poisson bracket (2.5), i.e. $\{\omega_n, \omega_m\} = 0$.

3 The normal form of conservation laws

Let us consider a formal conservation law of the form

$$u_t = \partial_x (u^2 + \epsilon a(u)u_x + \epsilon^2(b_1(u)u_{xx} + b_2(u)u_x^2) + O(\epsilon^3)). \quad (3.1)$$

It can be viewed as a higher order perturbation of the Hopf equation

$$u_t = \partial_x u^2 = 2uu_x. \quad (3.2)$$

Let us observe that there is no loss of generality by taking the Hopf equation as the leading order. Indeed, any equation of the form $u_t = K(u)u_x$ can be transformed into the equation (3.2) by a re-parametrization of the dependent variable $u = u(v)$. The Hopf equation (3.2) is Hamiltonian in the standard sense and completely integrable. Hence, there exist an infinite set of symmetries parametrized by a function of one variable $f(u)$ of the form

$$u_\tau = \partial_x f(u) = f'(u)u_x. \quad (3.3)$$

It is straightforward to check that $\partial_t u_\tau = \partial_\tau u_t$ for any function $f(u)$.

The aim of this section is to illustrate a perturbative approach to the problem of classifying integrable deformations of the Hopf hierarchy of the form (3.1). Various approaches have been developed to construct such deformations, and they can be briefly outlined as follows:

- *Symmetries approach*: It is based on the requirement that symmetry of the hydrodynamic limit can be extended to all perturbative orders [2, 33]. This is the most general approach because it does not require any additional assumption on the hierarchy. Nevertheless, it turns out to be computationally very expensive, in general.
- *Hamiltonian approach*: proposed by Dubrovin in [12] it consists in deforming Hamiltonian densities of the Hopf hierarchy with the request that deformed Hamiltonian functionals are still in involution.
- *bi-Hamiltonian approach*: proposed by Dubrovin and Zhang in [13] for the study of integrable PDEs appearing in the theory of Gromov-Witten invariants [24, 23] it is based on the extension of the bi-Hamiltonian structure to all perturbative orders.

The approach we propose here is a generalization of the Hamiltonian approach. Instead of deforming the Hamiltonian functionals we deform their differentials. Moreover, no assumption on the exactness of the deformed 1-forms is necessary. In this way, as mentioned in the previous section, we are able to treat simultaneously Hamiltonian and not Hamiltonian integrable deformations. Since the former have been already extensively studied in [12], we will mainly focus on the latter.

The classification procedure discussed in this paper is based on the following:

Definition 3.1 *A conservation law associated with the 1-form*

$$\omega_f^{def} = f(u) + \sum_{n=1}^{\infty} \epsilon^n g_n(u, u_x, \dots)$$

where each g_j is a homogenous differential polynomial of order j , is said to be integrable up to the order ϵ^k if there exists a 1-form

$$\omega_{\tilde{f}}^{def} = \tilde{f}(u) + \sum_{n=1}^{\infty} \epsilon^n \tilde{g}_n(u, u_x, \dots)$$

such that their Poisson bracket $\{\omega_f^{def}, \omega_{\tilde{f}}^{def}\} = 0$ vanishes identically modulo $O(\epsilon^{k+1})$ for any function $\tilde{f}(u)$.

Let us observe that taking for instance $f(u) = u^2$, the request that the two forms $\omega_{u^2}^{def}$ and $\omega_{\tilde{f}}^{def}$ commutes for any $\tilde{f}(u)$ up to a certain order in ϵ means that the corresponding deformed conservation law inherits all the symmetries of the Hopf equation up to the same order.

In the case of exact 1-forms the above commutativity condition reduces to the standard commutativity condition between Hamiltonian functionals with respect to a canonical Poisson bracket (see e.g. [12]).

A direct application of the definition (3.1) provides an effective tool to classify approximate integrable conservation laws viewed as deformations to the Hopf equation. The explicit derivation of integrability conditions becomes computationally more and more expensive as the order of such deformations increases. Nevertheless, it turns out that the most general deformation of the Hopf equation contains a certain number of redundant functional parameters that can be eliminated using the invariance of the form of conservation laws with respect to a special class of Miura transformations.

For instance, let us consider a general deformed conservation law of the form

$$u_t = \partial_x \omega_{u^2}^{def}(u, u_x, \dots; \epsilon) \quad (3.4)$$

with

$$\begin{aligned} \omega_{u^2}^{def} = & u^2 + \epsilon a(u)u_x + \epsilon^2 (b_1(u)u_{xx} + b_2(u)u_x^2) + \epsilon^3 (c_1(u)u_{xxx} + c_2(u)u_x u_{xx} + c_3(u)u_x^3) \\ & + \epsilon^4 (d_1(u)u_{4x} + d_2(u)u_x u_{xxx} + d_3(u)u_{xx}^2 + d_4(u)u_x^2 u_{xx} + d_5(u)u_x^4) + \epsilon^5 (e_1(u)u_{5x} \\ & + e_2(u)u_x u_{4x} + e_3(u)u_{xx} u_{xxx} + e_4(u)u_x^2 u_{xxx} + e_5(u)u_x u_{xx}^2 + e_6(u)u_x^3 u_{xx} + e_7(u)u_x^5) \\ & + O(\epsilon^6). \end{aligned}$$

We observe that the form of the equation (3.4) is preserved under the Miura transformation

$$u \rightarrow v = u + \epsilon^k \partial_x \beta(u, u_x, \dots) \quad (3.5)$$

where β is a homogeneous differential polynomial of degree $k - 1$. Hence, applying the Miura transformation (3.5) to the equation (3.4) we get

$$v_t = \partial_x (\omega_v(u(v, v_x, \dots), u_x(v, v_x, \dots), \dots) + \beta_t(u(v, v_x, \dots), u_x(v, v_x, \dots), \dots)),$$

where

$$u(v, v_x, \dots) = v - \epsilon^k \partial_x \beta(v, v_x, \dots) + O(\epsilon^{k+1}).$$

Clearly the equation truncated at the order $k - 1$ coincides with the original equation truncated at the same order. At the order k we have

$$\omega_k \rightarrow \tilde{\omega}_k = \omega_k(v, v_x, \dots) - 2v \partial_x \beta(v, v_x, \dots) + \sum_{s=0}^{k-1} \frac{\partial \beta(v, v_x, \dots)}{\partial v_{(s)}} \partial_x^{s+1} v^2$$

or equivalently

$$\tilde{\omega}_k = \omega_k(v, v_x, \dots) + \sum_{s=1}^{k-1} \frac{\partial \beta(v, v_x, \dots)}{\partial v_{(s)}} \left[\sum_{l=1}^s \binom{s+1}{l} v_{(l)} v_{(s+1-l)} \right]. \quad (3.6)$$

For instance, for $k = 2$ it is $\beta = \beta_1 v_x$ and

$$\tilde{\omega}_2 = b_1 v_{xx} + (b_2 + 2\beta_1) v_x^2.$$

Hence the term v_x^2 can be eliminated by choosing $\beta_1 = -b_2/2$. For $k = 3$ we have $\beta = \beta_{21} v_{xx} + \beta_{22} v_x^2$ and

$$\tilde{\omega}_3 = c_1(v) v_{xxx} + (c_2 + 6\beta_{21}) v_x v_{xx} + (c_3 + 4\beta_{22}) v_x^3.$$

Analogously, both terms $v_x v_{xx}$ and v_x^3 can be eliminated. Similarly, it is straightforward to check that for $k = 4$ one can retain just the terms v_{4x} and $v_x v_{xx}$, or alternatively v_{xx}^2 , while for $k = 5$ only v_{5x} and $v_x v_{4x}$, or alternatively $v_{xx} v_{xxx}$, will survive after a suitable Miura transformation.

The above examples suggest that it should be possible to choose β in such a way that only those terms that do not factor through v_x survive after the action of a suitable Miura transformation. This turns out to be true:

Theorem 3.2 *Given the conservation law*

$$u_t = \partial_x \omega_{u^2} \tag{3.7}$$

with $\omega_{u^2} = u^2 + \epsilon a(u) u_x + \sum_{k \geq 2} \epsilon^k \omega_k(u, u_x, \dots)$ there exists a sequence of Miura transformations of the form

$$u \rightarrow v = u + \epsilon^k \partial_x \beta(u, u_x, \dots)$$

that brings equation (3.7) into the normal form

$$v_t = \partial_x \omega_{v^2} \tag{3.8}$$

where

$$\omega_{v^2} = v^2 + \epsilon a(v) v_x + \sum_{k > 1} \epsilon^k \omega_k(v, v_x, \dots)$$

is such that

$$\frac{\partial \omega_k}{\partial v_x} = 0, \quad \forall k > 1.$$

Proof: Let us first introduce some useful notations. A general 1-form β homogeneous of degree $k - 1$ will be denoted as follows:

$$\sum_{i_1, \dots, i_{k-1} \quad i_1 + 2i_2 + \dots + (k-1)i_{k-1} = k-1} \beta_{[i_1, i_2, \dots, i_{k-1}]} v_{(1)}^{i_1} \dots v_{(k-1)}^{i_{k-1}}.$$

We will prove that the coefficients $\beta_{[i_1, i_2, \dots, i_{k-1}]}$ can be recursively determined through a lower triangular relation with respect to a suitable ordering of the monomials $v_{(1)}^{i_1} \dots v_{(k-1)}^{i_{k-1}}$. We introduce such an ordering, which is similar to a reverse

lexicographic ordering: the monomial $v_{(1)}^{i_1} \dots v_{(k-1)}^{i_{k-1}}$ ranks higher (or comes first or has a higher rank) than $v_{(1)}^{j_1} \dots v_{(k-1)}^{j_{k-1}}$, symbolically

$$v_{(1)}^{i_1} \dots v_{(k-1)}^{i_{k-1}} \succ v_{(1)}^{j_1} \dots v_{(k-1)}^{j_{k-1}},$$

if there exists $m \in \{1, \dots, k-1\}$ such that $i_l = j_l$ for all $l > m$ and $i_m > j_m$. In other words, monomials are ranked according to the higher derivative of v . For instance, among the homogenous monomials of degree $k-1$, the highest ranking monomial is $v_{(k-1)}$ and the lowest ranking is $v_{(1)}^{k-1}$. This gives clearly a total ordering of the monomials. In the following, we will write down a 1-form homogeneous of a certain degree from the highest to the lowest ranking. For instance, for a homogeneous 1-form of degree 3

$$\beta = \beta_{[0,0,1]} v_{(3)} + \beta_{[1,1,0]} v_{(1)} v_{(2)} + \beta_{[3,0,0]} v_{(1)}^3.$$

Let us now fix k and write ω_k as $\alpha + v_{(1)}\eta$ where η is ordered as above. In the rest of the proof α will be neglected as we will focus only on the terms of $v_{(1)}\eta$ that factors through $v_{(1)}$.

Let us now expand the expression

$$\sum_{s=1}^{k-1} \frac{\partial \beta(v, v_x, \dots)}{\partial v_{(s)}} \left[\sum_{l=1}^s \binom{s+1}{l} v_{(l)} v_{(s+1-l)} \right] \quad (3.9)$$

entering the Miura transformation (3.6) as follows

$$\begin{aligned} \sum_{s=1}^{k-1} \frac{\partial \beta(v, v_x, \dots)}{\partial v_{(s)}} 2(s+1) v_{(s)} v_{(1)} + \sum_{s=3}^{k-1} \frac{\partial \beta(v, v_x, \dots)}{\partial v_{(s)}} \left[\sum_{l=2}^{s-1} \binom{s+1}{l} v_{(l)} v_{(s+1-l)} \right] \\ = \tilde{\beta} v_{(1)} + \delta, \end{aligned}$$

where

$$\tilde{\beta} = \sum_{s=1}^{k-1} \frac{\partial \beta(v, v_x, \dots)}{\partial v_{(s)}} 2(s+1) v_{(s)}$$

is meant to be ordered as above and the residual term is

$$\delta = \sum_{s=3}^{k-1} \frac{\partial \beta(v, v_x, \dots)}{\partial v_{(s)}} \left[\sum_{l=2}^{s-1} \binom{s+1}{l} v_{(l)} v_{(s+1-l)} \right]. \quad (3.10)$$

Notice that the coefficients in $\tilde{\beta}$ coincide, up to a positive factor, with the coefficients in β , that is

$$\tilde{\beta}_{[i_1, i_2, \dots, i_{k-1}]} = c_{i_1, \dots, i_{k-1}} \beta_{[i_1, \dots, i_{k-1}]},$$

where $c_{i_1, \dots, i_{k-1}}$ are positive constants. In particular, terms in $\tilde{\beta}$ have the same ranking of the corresponding terms in β .

We are now able to prove that it is always possible to eliminate all terms containing $v_{(1)}$ starting from the highest ranking coefficients. However, special attention should be paid as one needs to take into account contributions coming from both $\tilde{\beta}v_{(1)}$ and $\eta v_{(1)}$ and δ . It will turn out that all such contributions appear in a lower triangular form and then we can recursively fix coefficients of $\tilde{\beta}$ and consequently β in such a way to kill $\eta v_{(1)}$.

Let us first note that it is always possible to fix the highest ranking coefficient in $\tilde{\beta}v_{(1)}$ that annihilates the highest ranking coefficient in $v_{(1)}\eta$. Indeed, the highest ranking term in $\tilde{\beta}v_{(1)}$ is $\tilde{\beta}_{[0,0,\dots,1]}v_{(1)}v_{(k-1)}$ while the highest ranking term in $\eta v_{(1)}$ is $\eta_{[0,0,\dots,1]}v_{(1)}v_{(k-1)}$. It is immediate to check that no such a term can originate from δ and therefore $\tilde{\beta}_{[0,0,\dots,1]}$ is uniquely determined by the condition $\tilde{\beta}_{[0,0,\dots,1]} + \eta_{[0,0,\dots,1]} = 0$.

Let us now proceed by induction following the ordering (from the highest to the lowest ranking) of the monomials. Assuming that the L highest ranking monomial in $\tilde{\beta}$ is determined, we show that it is possible to find the $L+1$ highest ranking monomial $\tilde{\beta}_{[i_1,\dots,i_{k-1}]}v_{(1)}^{i_1+1} \dots v_{(k-1)}^{i_{k-1}}$ in $\tilde{\beta}$ that annihilates the corresponding $L+1$ highest ranking monomial $\eta_{[i_1,\dots,i_{k-1}]}v_{(1)}^{i_1+1} \dots v_{(k-1)}^{i_{k-1}}$ in $\eta v_{(1)}$ by imposing the condition

$$\eta_{[i_1,\dots,i_{k-1}]}v_{(1)}^{i_1+1} \dots v_{(k-1)}^{i_{k-1}} + \tilde{\beta}_{[i_1,\dots,i_{k-1}]}v_{(1)}^{i_1+1} \dots v_{(k-1)}^{i_{k-1}} + \text{residual terms} = 0, \quad (3.11)$$

where residual terms come from δ . Indices $i_1, \dots, i_{k-1}$ above are fixed.

From its definition, δ can be thought as an operator acting linearly on β , that is $\delta(\beta_1 + \beta_2) = \delta(\beta_1) + \delta(\beta_2)$. Therefore, without loss of generality, we can focus our attention on the action of δ on the generic term of the form

$$\gamma = \beta_{[j_1,\dots,j_{k-1}]}v_{(1)}^{j_1} \dots v_{(k-1)}^{j_{k-1}}.$$

Up to a positive factor, we can write

$$\delta(\gamma) = \beta_{[j_1,\dots,j_{k-1}]} \sum_{s=3}^{k-1} \sum_{l=2}^{s-1} v_{(1)}^{j_1} \dots v_{(l)}^{j_l+1} \dots v_{(s+1-l)}^{j_{s+1-l}+1} \dots v_{(s)}^{j_s-1} \dots v_{(k-1)}^{j_{k-1}}. \quad (3.12)$$

Since $l \leq s-1 < s$ and analogously $s+1-l < s$ and j_s is mapped into j_s-1 , from equation (3.12) the operator δ always decreases the ranking of the terms on which it acts.

This means that the residual term into the equation (3.11) only contains coefficients in β (i.e. $\tilde{\beta}$) already determined when considering the corresponding equation at the higher orders. Hence we can solve the linear equation (3.11) w.r.t $\tilde{\beta}_{[i_1,\dots,i_{k-1}]}$ in terms of $\eta_{[i_1,\dots,i_{k-1}]}$ and in terms of already determined higher ranking coefficients. This concludes our proof. ■

4 Classification results

We now proceed to the classification of involutive 1-forms

$$\begin{aligned}\omega_{u^2}^{def} = & u^2 + \epsilon a(u)u_x + \epsilon^2 b_1(u)u_{xx} + \epsilon^3 c_1(u)u_{xxx} + \epsilon^4 [d_1(u)u_{4x} + d_2(u)u_{xx}^2] + \\ & + \epsilon^5 [e_1(u)u_{5x} + e_2(u)u_{xx}u_{xxx}] + \dots\end{aligned}\quad (4.1)$$

$$\begin{aligned}\omega_f^{def} = & f(u) + \epsilon Au_x + \epsilon^2 (B_1u_{xx} + B_2u_x^2) + \epsilon^3 (C_1u_{xxx} + C_2u_xu_{xx} + C_3u_x^3) \\ & + \epsilon^4 (D_1u_{4x} + D_2u_xu_{xxx} + D_3u_{xx}^2 + D_4u_x^2u_{xx} + D_5u_x^4) \\ & + \epsilon^5 (E_1u_{5x} + E_2u_xu_{4x} + E_3u_{xx}u_{xxx} + E_4u_x^2u_{xxx} + E_5u_xu_{xx}^2 + E_6u_x^3u_{xx} + E_7u_x^5) \\ & + \dots\end{aligned}\quad (4.2)$$

up to a certain order ϵ^k w.r.t. the Poisson bracket defined in (2.5), that is

$$\{\omega_{u^2}^{def}, \omega_f^{def}\} = O(\epsilon^{k+1}).$$

In virtue of the theorem (3.2) we can take $\omega_{u^2}^{def}$ in its normal form (4.1) without loss of generality. We have the following

Theorem 4.1 *If $a(u) \neq 0$, then up to $O(\epsilon^6)$, the normal forms of integrable conservation laws*

$$u_t = \partial_x \omega_{u^2}^{def}$$

and their commuting flows

$$u_\tau = \partial_x \omega_f^{def}$$

are parametrized only by the functional parameter $a(u)$ (here $\omega_{u^2}^{def}$ and ω_f^{def} are respectively given by (4.1) and (4.2)).

Proof: The constraint that the Poisson bracket $\{\omega_{u^2}^{def}, \omega_f^{def}\}$ vanishes up to the order $O(\epsilon^6)$ provides the following set of conditions on the deformation coefficients

$O(\epsilon^0) \rightarrow$ no conditions

$O(\epsilon) \rightarrow$ Gives A in terms of a and f :

$$A(u) = \frac{1}{2}a(u)f''(u) \quad (4.3)$$

$O(\epsilon^2) \rightarrow$ Gives B_1 and B_2 in terms of b_1 , f and $a(u)$:

$$\begin{aligned}
B_1 &= \frac{1}{2}b_1f'' + \frac{1}{6}a^2f''' \\
B_2 &= \frac{1}{4}aa'f''' + \frac{1}{8}a^2f^{(4)} + \frac{1}{4}b_1f'''
\end{aligned}$$

$O(\epsilon^3) \rightarrow$ Gives C_1 , C_2 and C_3 in terms of the small letters and f

$$\begin{aligned}
C_1 &= \frac{1}{3}a^2a'f''' + \frac{1}{2}c_1f'' + \frac{1}{24}a^3f^{(4)} \\
C_2 &= \frac{11}{12}a(a')^2f''' + \frac{5}{6}a^2a'f^{(4)} + \frac{7}{24}a^2a''f''' + \frac{3}{4}c_1f''' + \frac{1}{12}a^3f^{(5)} \\
C_3 &= \frac{1}{3}aa'a''f''' + \frac{11}{24}a(a')^2f^{(4)} + \frac{1}{6}c_1f^{(4)} + \frac{1}{48}a^3f^{(6)} + \frac{1}{18}a^2a'''f''' + \frac{1}{6}a^2a''f^{(4)} + \frac{1}{4}a^2a'f^{(5)}
\end{aligned}$$

together with the constraint

$$b_1 = (a^2/2!)' \quad (4.4)$$

$O(\epsilon^4) \rightarrow$ Gives D_1 , D_2 , D_3 , D_4 and D_5 in terms of the small letters and f

$$\begin{aligned}
D_1 &= \frac{1}{8}a^3a'f^{(4)} + \frac{1}{6}a^3a''f''' + \frac{1}{120}a^{(4)}f^{(5)} + \frac{1}{2}a^2(a')^2f''' + \frac{1}{2}d_1f'' \\
D_2 &= \frac{9}{16}a^3a''f^{(4)} + \frac{1}{2}d_2a'f'' + \frac{7}{4}a^2a'a''f''' + d_1f''' + \frac{1}{6}a^3a'''f''' + \frac{1}{48}a^4f^{(6)} + \frac{3}{8}a^3a'f^{(5)} + \\
&\quad + \frac{15}{8}a^2(a')^2f^{(4)} + \frac{3}{2}a(a')^3f''' \\
D_3 &= \frac{17}{24}a^2a'a''f''' + \frac{1}{72}a^3a'''f''' + \frac{17}{48}a^3a''f^{(4)} + \frac{11}{12}a(a')^3f''' + \frac{5}{4}a^2(a')^2f^{(4)} + \frac{3}{4}d_1f''' + \\
&\quad + \frac{1}{72}a^4f^{(6)} + \frac{1}{4}a^3a'f^{(5)} \\
D_4 &= \frac{7}{16}a^3a'2f^{(6)} + \frac{3}{4}a^3a''f^{(5)} + \frac{29}{30}a^2a'a'''f''' + \frac{27}{8}a(a')^3f^{(4)} + \frac{1}{48}a^4f^{(7)} + \frac{3}{5}d_2f''' + \\
&\quad + \frac{29}{10}a(a')^2a''f''' + 4a^2a'a''f^{(4)} + d_1f^{(4)} + \frac{21}{8}a^2(a')^2f^{(5)} + \frac{1}{3}a^3a'''f^{(4)} + \\
&\quad + \frac{1}{12}a^3a'a^{(4)}f''' + \frac{9}{10}a^2(a'')^2f''' \\
D_5 &= \frac{23}{576}a^3a^{(4)}f^{(4)} + \frac{1}{144}a^3a^{(5)}f''' + \frac{19}{48}a^2(a'')^2f^{(4)} + \frac{1}{8}d_2f^{(4)} + \frac{1}{8}a^3a''f^{(6)} + \frac{13}{144}a^3a'''f^{(5)} + \\
&\quad + \frac{3}{4}a(a')^3f^{(5)} + \frac{1}{384}a^4f^{(8)} + \frac{23}{144}a^2a''a'''f''' + \frac{7}{16}a^2(a')^2f^{(6)} + \frac{3}{16}aa'(a'')^2f''' + \\
&\quad + \frac{1}{16}a^3a'f^{(7)} + \frac{1}{8}d_1f^{(5)} + \frac{73}{144}a^2a'a'''f^{(4)} + \frac{13}{144}a^2a'a^{(4)}f''' + \frac{47}{48}a^2a'a''f^{(5)} + \\
&\quad + \frac{7}{18}a(a')^2a'''f''' + \frac{4}{3}a(a')^2a''f^{(4)}.
\end{aligned}$$

and the constraint

$$c_1 = (a^3/3!)'' \quad (4.5)$$

$O(\epsilon^5) \rightarrow$ Gives $E_1, E_2, E_3, E_4, E_5, E_6$ and E_7 in terms of the small letters and f (whose cumbersome expression is not presented here) and the constraints

$$\begin{aligned} d_1 &= (a^4/4!)''' \\ d_2 &= \frac{5}{24}a^3a^{(4)} + \frac{8}{3}a(a')^2a'' + a^2(a'')^2 + \frac{31}{18}a^2a'a''' . \end{aligned}$$

■

We call *integrable viscous conservations laws* all conservation laws obtained from this deformation procedure extended at any order in the parameter ϵ and such that $a(u) \neq 0$. The perturbative calculations performed above support the formulation of the following

Conjecture 4.2 *The normal form of integrable viscous conservation laws associated with the 1-form (4.1) as well as their commuting flows associated with the 1-form (4.2) are uniquely determined at any order in ϵ by the non-vanishing functional parameter $a(u)$.*

It should also be noted that if $a(u)$ is allowed to be vanishing the deformation procedure develops a branching. The branch $a(u) = 0$ corresponds to the case of dispersive perturbations. In particular, Hamiltonian dispersive perturbations have been extensively studied in a number of papers (see e.g. [13, 12, 23]). It is conjectured that all deformations are uniquely specified by a number of functional parameters that in the bi-Hamiltonian setting are called central invariants [14]. The functional parameter $a(u)$ plays in the present context the same role as central invariants in the Hamiltonian setup. Hence, we refer to $a(u)$ as *viscous central invariant*.

An important well-known example of integrable viscous conservation law as been already mentioned above and it is given by the Burgers equation

$$u_t = \partial_x (u^2 + \epsilon u_x) = 2uu_x + \epsilon u_{xx}, \quad (4.6)$$

Let us observe that Burgers' equation already appears in its normal form and it possesses a constant viscous central invariant $a(u) = 1$. Conversely, one can immediately check that in the case of constant viscous central invariant the 1-form $\omega_{u^2}^{def}$ truncates at the first order in ϵ providing the Burgers equation. Higher flows of the Burgers hierarchy can be obtained as a deformation of higher flows of the Hopf hierarchy obtained choosing in (4.2) $f(u) = u^n$, $n \geq 2$.

5 Linear viscous central invariants

5.1 The viscous analogue of the Camassa-Holm equation

In the present section we show that the deformation procedure discussed above provides, in the case of the linear viscous central invariant $a(u) = u$, the following

non-evolutionary conservation law

$$u_t - \epsilon u_{xt} = \partial_x (u^2 - \epsilon u u_x). \quad (5.1)$$

Equation (5.1) first appeared as a scalar reduction of two-component Camassa-Holm type equations [31, 16, 22].

This equation belongs to the class of viscous conservation laws under consideration and can be written in evolutionary form as a formal series in ϵ . In fact, the application of the formal inverse operator

$$(1 - \epsilon \partial_x)^{-1} = 1 + \epsilon \partial_x + \epsilon^2 \partial_x^2 + \dots = \sum_{k=0}^{\infty} \epsilon^k \partial_x^k \quad (5.2)$$

to both sides of (5.1) gives

$$\begin{aligned} u_t &= (1 - \epsilon \partial_x)^{-1} [2uu_x - \epsilon(uu_{xx} + u_x^2)] \\ &= (1 - \epsilon \partial_x)^{-1} \partial_x \left[\frac{u^2}{2} + (1 - \epsilon \partial_x) \left(\frac{u^2}{2} \right) \right] \\ &= \partial_x \left[\frac{u^2}{2} + \sum_{k=0}^{\infty} \epsilon^k \partial_x^k \left(\frac{u^2}{2} \right) \right]. \end{aligned} \quad (5.3)$$

We show that equation (5.1), or equivalently (5.3), can be brought to the normal form via a Miura transformation. This statement is made precise by the following

Theorem 5.1 *The equation (5.1) is reduced to its normal form*

$$u_t = \partial_x \left(\sum_k \epsilon^k u u_{(k)} \right) \quad (5.4)$$

via the Miura transformation

$$u = v - \epsilon v_x. \quad (5.5)$$

where $v = v(x, t)$ is a solution to the equation (5.1) or, equivalently, (5.3).

Proof: Inverting the Miura transformation (5.5) we have

$$v = (1 - \epsilon \partial_x)^{-1} u.$$

Differentiating by t and using the equation (5.4) we get

$$\begin{aligned} v_t &= (1 - \epsilon \partial_x)^{-1} u_t = (1 - \epsilon \partial_x)^{-1} \partial_x \left(\sum_{k=0}^{\infty} \epsilon^k u u_{(k)} \right) = \partial_x (1 - \epsilon \partial_x)^{-1} u v \\ &= \partial_x (1 - \epsilon \partial_x)^{-1} (v^2 - \epsilon v v_x) = \partial_x \sum_{k=0}^{\infty} \epsilon^k \partial_x^k \left[v^2 - \epsilon \partial_x \left(\frac{v^2}{2} \right) \right] \\ &= \partial_x \left[v^2 + \sum_{k=1}^{\infty} \epsilon^k \partial_x^k \left(\frac{v^2}{2} \right) \right] = \partial_x \left[\frac{v^2}{2} + \sum_{k=0}^{\infty} \epsilon^k \partial_x^k \left(\frac{v^2}{2} \right) \right] \end{aligned}$$

where we used the expansion formula (5.2). The theorem is proved. ■

Comparing with the results of the previous section we see that up to the fifth order in the deformation parameter the equation (5.4) coincides with the equation

$$u_t = \partial_x \omega_{u^2}^{def}$$

with $a(u) = u$. According with the Conjecture 4.2, the equation (5.4) is expected to be the only scalar integrable viscous conservation law, up to Miura transformations, possessing a linear viscous central invariant. We note that the positive hierarchy is obtained as the set of symmetries corresponding to the choice $f(u) = u^n$, $n = 0, 1, 2, 3, \dots$ (including the original equation) while the negative hierarchy represents the set of symmetries corresponding to the choice $f(u) = u^{-n}$, $n = 1, 2, 3, \dots$. In the next subsection we write explicitly the negative and positive hierarchies and provide an alternative proof of the integrability of equation (5.1) via linearization.

5.2 The negative hierarchy

The analysis of deformations discussed in the Section 4 suggests that the normal form of the equation (5.1) is not truncated. Nevertheless, there exists a family of flows in the hierarchy possessing a finite deformation, that is the negative hierarchy

$$u_{t_{-n}} = \partial_x \left(\frac{1}{u} - \epsilon \partial_x \frac{1}{u} \right)^n (1) \quad n = 1, 2, 3, \dots \quad (5.6)$$

The first equation of this hierarchy obtained for $n = 1$

$$u_{t_{-1}} = \partial_x \left(\frac{1}{u} - \epsilon \partial_x \frac{1}{u} \right). \quad (5.7)$$

has been included by F. Calogero [4] within a list of nonlinear PDEs that are integrable via a nonlinear change of variables. Subsequently, it was observed in [31] that the equation (5.7) can be mapped into Burgers' equation. Let us note that the hierarchy (5.6) can be written equivalently in terms of a recursion operator as

$$u_{t_{-n}} = R^{-1} u_{t_{-1}}$$

where

$$R^{-1} = \partial_x (1 - \epsilon \partial_x) \frac{1}{u} \partial_x^{-1} \quad (5.8)$$

is the inverse of the operator

$$R = \partial_x u (1 - \epsilon \partial_x)^{-1} \partial_x^{-1} \quad (5.9)$$

such that equation (5.4) can be written as

$$u_t = R u_x. \quad (5.10)$$

Operators (5.8) and (5.9) can be viewed as a reduction of the recursion operators for, respectively, negative and positive hierarchies of the system studied in [16]. A direct proof that (5.8) and (5.9) are indeed recursion operators is based on the application on a criterion proposed by P. Olver in [30].

We now prove the integrability of equation (5.1) in two steps. First, we show that the flows (5.6) can be mapped to the heat hierarchy by a sequence of reciprocal and nonlinear transformations and as a consequence commute. Secondly, we prove that the equation (5.4) transforms into the Klein-Gordon equation and consequently commutes with the flows (5.6).

Lemma 5.2 *The family of flows (5.6) is transformed into the Burgers hierarchy via the reciprocal transformation*

$$v(\varphi, t_{-1}, \dots, t_{-n}) = -\partial_\varphi x(\varphi, t_{-1}, \dots, t_{-n}) \quad \text{where} \quad \varphi = \int u \, dx, \quad (5.11)$$

that is the function v is a simultaneous solution to the Burgers hierarchy

$$v_{t_{-n}} = \partial_\varphi (v + \epsilon \partial_\varphi)^n v \quad n = 1, 2, 3, \dots$$

Proof Introducing the potential φ such that $u = \varphi_x$, equations (5.6) after the integration w.r.t. x give

$$\varphi_{t_{-n}} = \left(\frac{1}{\varphi_x} - \epsilon \partial_x \frac{1}{\varphi_x} \right)^n (1). \quad (5.12)$$

For $n = 1$ the above equation takes the form

$$\varphi_{t_{-1}} = \frac{1}{\varphi_x} + \epsilon \frac{\varphi_{xx}}{\varphi_x^2}. \quad (5.13)$$

Introducing the reciprocal transformation of the form $x = x(\varphi, t_1, t_2, \dots)$, one has

$$\varphi_x = \frac{1}{x_\varphi} \quad \varphi_{xx} = -\frac{x_{\varphi\varphi}}{x_\varphi^3} \quad \varphi_{t_n} = -\frac{x_{t_n}}{x_\varphi}, \quad n = 1, 2, \dots \quad (5.14)$$

Using relations above into the equation (5.13) one arrives to

$$x_{t_{-1}} = (-x_\varphi + \epsilon \partial_\varphi) x_\varphi$$

that is just the potential Burgers equation, i.e. $v = -x_\varphi$ satisfies the Burgers equation

$$v_{t_{-1}} = \partial_\varphi (v + \epsilon \partial_\varphi) v.$$

Let us now proceed by induction assuming that the proposition is true for the n -th equation and consider the $(n+1)$ -th equation

$$\varphi_{t_{-n-1}} = \left(\frac{1}{\varphi_x} - \epsilon \partial_x \frac{1}{\varphi_x} \right)^{n+1} (1) = \left(\frac{1}{\varphi_x} - \epsilon \partial_x \frac{1}{\varphi_x} \right) \varphi_{t_{-n}}.$$

Using the relations (5.14) and observing that $\partial_x = x_\varphi^{-1} \partial_\varphi$, we arrive to the equation

$$x_{t_{-n-1}} = (x_\varphi - \epsilon \partial_\varphi) x_{t_{-n}}.$$

Since

$$x_{t_{-n}} = (-x_\varphi + \epsilon \partial_\varphi)^n x_\varphi$$

we have

$$x_{t_{-n-1}} = -(-x_\varphi + \epsilon \partial_\varphi)^{n+1} x_\varphi.$$

Finally, let us differentiate the expression above by φ and substitute $x_\varphi = -v$. The lemma is proved. $\blacksquare$

Theorem 5.3 *The flows (5.6) commute.*

Proof: In virtue of the Lemma (5.2) the set of flows (5.6) is transformed into Burgers' hierarchy via the reciprocal transformation (5.11). On the other hand it is known that the Cole-Hopf transformation $v = \epsilon \partial_\varphi \log w$ brings Burgers' hierarchy into the heat hierarchy

$$w_{t_{-n}} = \epsilon^n \partial_\varphi^{n+1} w.$$

This result can be straightforwardly verified for $n = 1$ and then proved by induction for any n . Flows of the heat hierarchy clearly commute, i.e. $\partial_{t_{-n}} \partial_{t_{-m}} w = \partial_{t_{-m}} \partial_{t_{-n}} w = \epsilon^{n+m} \partial_\varphi^{n+m+2} w$. The theorem is proved. $\blacksquare$

The following theorem uncovers the relation existing between the equation (5.1) presented above and the integrable hierarchy (5.6).

Theorem 5.4 *The flow of conservation law (5.4) commutes with all flows of the hierarchy (5.6).*

Proof: Let us note that the equation (5.4) can be written in the more compact form

$$v_t = \partial_x [v(1 - \epsilon \partial_x)^{-1} v]. \quad (5.15)$$

Introducing the variable φ defined by $\varphi_x = v$, integration with respect to x provides us with the equation

$$\varphi_t \varphi_x - \varphi_x^3 + \epsilon (\varphi_t \varphi_{xx} - \varphi_x \varphi_{xt}) = 0, \quad (5.16)$$

where we used the fact that the integration constant can always be eliminated by a shift of the independent variable $\varphi \rightarrow \varphi + f(t)$. The reciprocal transformation $x = x(\varphi, t)$ brings the equation above into the form

$$\epsilon x_{\varphi t} - x_\varphi x_t = 1,$$

and the change of variable $x = -\epsilon \log w$ maps the equation above into the Klein-Gordon equation in light cone variables

$$\epsilon^2 w_{\varphi t} + w = 0.$$

Note that both the hierarchy (5.6) and the equation (5.4) can be linearized via the same change of variables that brings them to the heat hierarchy and Klein-Gordon equation respectively. Since the Klein-Gordon equation is compatible with all members of the heat hierarchy, the theorem is proved. $\blacksquare$

5.3 The positive hierarchy

Given the equation (5.4) in the form (5.10) it is natural to consider the family of flows of the form

$$u_{t_n} = R^n u_x, \quad n = 1, 2, \dots \quad (5.17)$$

where R is defined in (5.9). We now prove that the family of flows (5.17) can be mapped into a family of linear commuting flows whose first member is the Klein-Gordon equation and for this reason we name it *Klein-Gordon hierarchy*. We have the following

Theorem 5.5 *The family of flows (5.17) is mapped to the Klein-Gordon hierarchy that is given by the following set of linear commuting flows*

$$w_{t_n} = \epsilon^{-n} \partial_{\varphi}^{-n} w, \quad n = 1, 2, \dots \quad (5.18)$$

via the nonlinear change of variables

$$x(\varphi, t_1, t_2, \dots) = -\epsilon \log w$$

where

$$x = \varphi^{-1} \quad \text{and} \quad \varphi = \int u \, dx.$$

Moreover, the positive hierarchy (5.17) and the negative hierarchy (5.6) commute.

Proof: Let us first note that the theorem holds true for $n = 1$ in virtue of the proof of the theorem (5.4). Let us now consider the equations (5.17) for $n \geq 2$ in the form

$$u_{t_n} = R u_{t_{n-1}}, \quad n = 1, 2, \dots$$

Introducing the function φ such that $u = \varphi_x$, following the same steps as in the $n = 1$ case the above equation can be equivalently written as

$$\varphi_x \varphi_{t_n} + \epsilon (\varphi_{xx} \varphi_{t_n} - \varphi_x \varphi_{x t_n}) = \varphi_x^2 \varphi_{t_{n-1}}.$$

Introducing the reciprocal transformation

$$x = x(\varphi, t_1, t_2, \dots)$$

we arrive to the equation

$$\epsilon x_{\varphi t_n} - x_{\varphi} x_{t_n} + x_{t_{n-1}} = 0. \quad (5.19)$$

Setting $x = -\epsilon \log w$ one readily gets the following linear equations

$$\epsilon w_{\varphi t_n} + w_{t_{n-1}} = 0, \quad n = 2, 3, \dots$$

that can be equivalently written in the evolutionary non-local form (5.18). It is immediate to check the commutativity condition for the flows of both the positive and the negative hierarchy, i.e. $\partial_{t_n} \partial_{t_m} w = \partial_{t_m} \partial_{t_n} w$ for any $n, m \in \mathbb{Z}$. The theorem is proved. $\blacksquare$

Summarizing, the normal form of equations associated with a linear viscous central invariant can be mapped into the Burgers equation by means of a reciprocal transformation and then to the heat equation via a Cole-Hopf transformation. Interestingly, the same sequence of transformations brings on one hand the negative hierarchy into the *positive* Burgers hierarchy and then to the heat hierarchy. On the other hand, the positive hierarchy is mapped into the *negative* Burgers hierarchy whose potential form is given by the recursive nonlocal equations (5.19) that is linearised into the Klein-Gordon hierarchy (see also Figure 1)). Hence, the Klein-Gordon hierarchy can be interpreted as the negative heat hierarchy. Indeed, introducing the recursion operator

$$\mathcal{R} = \epsilon \partial_{\varphi}$$

the Burgers' hierarchy is written as

$$w_{t_{-n}} = \mathcal{R}^n w_{\varphi} \quad n = 0, 1, 2, \dots$$

while the Klein-Gordon hierarchy is

$$w_{t_n} = \mathcal{R}^{-n} w \quad n = 1, 2, \dots$$

6 Local well-posedness of the Cauchy problem

6.1 Kato's method

The present Section is devoted to the study of the local (in time) well-posedness of the periodic Cauchy problem for (5.1) by using Kato's method [18] that is here briefly outlined.




Figure 1: Positive and negative integrable hierarchies associated with viscous conservation laws possessing a linear viscous central invariant.

Let us consider the Cauchy problem for the quasi-linear equation of evolution:

$$\frac{d}{dt}v + A(v)v = f(v), \quad t > 0 \quad v(0) = \phi. \quad (6.1)$$

Let X and Y be reflexive Banach spaces with Y continuously and densely embedded in X , let $S : Y \rightarrow X$ be an isomorphism and suppose that the norm of Y is chosen in such a way that S is an isometry. Assume now that the terms appearing in (6.1) satisfy the following conditions:

1. A is an operator-valued function defined on Y , that is for each $y \in Y$, $A(y) : D(A) \subset X \rightarrow X$ is a linear operator on X (in the interesting cases $A(y)$ is an unbounded operator). Assume that $A(y)$ is quasi-m-accretive, uniformly for $y \in Y$ with $\|y\|_Y \leq M$. In other words, for every constant $M > 0$, there is a real number ω such that for every $y \in Y$, with $\|y\|_Y \leq M$, $-A(y)$ generates a C^0 -semigroup $\{e^{-tA(y)}\}_{t \geq 0}$ with

$$\|e^{-tA(y)}\|_{\mathcal{L}(X)} \leq e^{\omega t}, \quad t \geq 0,$$

where $\|\cdot\|_{\mathcal{L}(X)}$ is the operator norm.

2. For each $y \in Y$, $A(y)$ is a bounded linear operator from Y to X and moreover

$$\|(A(y) - A(z))w\|_X \leq \mu_A \|y - z\|_X \|w\|_Y, \quad y, z, w \in Y, \quad (6.2)$$

for some constant μ_A depending only on $\max\{\|y\|_Y, \|z\|_Y\}$.

3. for any $M > 0$, the inequality

$$\|(SA(y) - A(y)S)S^{-1}w\|_X \leq \mu_1(M) \|w\|_X, \quad y \in Y, \|y\|_Y \leq M \quad (6.3)$$

holds for all $w \in Y$, where $\mu_1(M) > 0$ is a constant.

4. For each $M > 0$, f is a bounded function from $\{y \in Y : \|y\|_Y \leq M\}$ to Y . Also we have

$$\|f(y) - f(z)\|_X \leq \mu_2 \|y - z\|_X, \quad y, z \in Y, \quad (6.4)$$

and

$$\|f(y) - f(z)\|_Y \leq \mu_3 \|y - z\|_Y, \quad y, z \in Y, \quad (6.5)$$

for some constants μ_2 and μ_3 , where μ_2 depends only on $\max\{\|y\|_Z, \|z\|_Z\}$ and μ_3 depends only on $\max\{\|y\|_Y, \|z\|_Y\}$, where Z is a space such that $Y \subset Z \subset X$, all with continuous inclusions. (In our application it will be $X := L^2, Y := H^1, Z := L^\infty$ all for functions defined on the circle.)¹

Then one has the following Theorem (see [18]):

Theorem 6.1 *Assume conditions (1), (2), (3) and (4) above hold. Then for any $\phi \in Y$, there is a $T > 0$, depending only on $\|\phi\|_Y$ and a unique solution v to (6.1) such that $v \in C^0([0, T], Y) \cap C^1([0, T], X)$. Moreover, $v(t)$ depends continuously on the initial data $\phi = v(0)$ in the Y -norm.*

6.2 The periodic Cauchy problem for the equation (5.1)

Let us consider the equation (5.1) with fixed ϵ and, for convenience, we set $\epsilon = 1$ and replace $t \rightarrow -t$. With these notations the initial value problem for the equation (5.1) will read as follows

$$m_t = -vm_x - v_x m \quad v(x, 0) = w(x) \quad (6.6)$$

where $m := v - v_x$ and $w : \mathbb{S}^1 \rightarrow \mathbb{R}$. The above time inversion will result into proving the local well-posedness of the Cauchy problem for equation (5.1) backward in time. We will finally observe that the result holds also true for the forward evolution.

Using the fact that $v = S^{-1}m$ where $S = 1 - \partial_x$ the problem (6.6) can be equivalently formulated as follows

$$m_t = -(S^{-1}m)m_x - m(S^{-1}m)_x, \quad m(x, 0) = \phi(x) \quad (6.7)$$

where $\phi : \mathbb{S}^1 \rightarrow \mathbb{R}$. Verifying the conditions for the validity of Kato's theorem will guarantee well-posedness for $\phi \in H^1(\mathbb{S}^1)$.

From now on all the functional spaces introduced are referred to functions on the circle and we for sake of simplicity will be denoted just as L^2, H^1 .

We have the following

¹Note that these assumptions are exactly like those stated in [7], except for the Lipschitz inequality (6.4). Indeed in [7] it is required that μ_2 depends only on $\max\{\|y\|_X, \|z\|_X\}$, while here we require dependence on a bigger quantity. As it can be seen in [18] page 40, second inequality from above, this is not affecting the statement of Theorem 6.1 in our case.

Theorem 6.2 *Let $\phi \in H^1$, then there is a $T > 0$, depending only on $\|\phi\|_{H^1}$ such that (6.7) has a unique solution $m \in C^0([0, T], H^1) \cap C^1([0, T], L^2)$ and moreover m as a function of t depends continuously on ϕ in the H^1 -norm.*

Proof: We have just to check assumptions (1) to (4) of Theorem 6.1. We choose as $X = L^2$ and as $Y = H^1$. On L^2 we use the usual L^2 -norm. Recall that H^1 is continuously and densely embedded in L^2 , since for any $y \in H^1$, $\|y\|_{L^2} \leq \|y\|_{H^1}$ (by definition $\|y\|_{H^1}^2 := \|y\|_{L^2}^2 + \|y_x\|_{L^2}^2$). Also we choose as an isometric isomorphism $S = 1 - \partial_x$, so that

$$\begin{aligned} \|Sy\|_X^2 &= \|y - y_x\|_{L^2}^2 = \langle y - y_x, y - y_x \rangle_{L^2} = \\ &= \|y\|_{L^2}^2 + \|y_x\|_{L^2}^2 - \langle y_x, y \rangle_{L^2} - \langle y, y_x \rangle_{L^2}, \end{aligned}$$

where the terms $\langle y_x, y \rangle_{L^2} + \langle y, y_x \rangle_{L^2}$ give a zero contribution due to integration by parts which holds for functions in H^1 (see for instance Corollary 8.1 [3]). Therefore, with the choice of S as an isometric isomorphism, the norm on H^1 is equal to the standard norm on H^1 . Comparing (6.7) with (6.1), we choose

$$A(y) = (S^{-1}y)\partial_x, \quad f(y) = -y(S^{-1}y)_x, \quad y \in Y = H^1.$$

We start verifying assumption (1), fixing $M > 0$ and $y \in Y$ with $\|y\|_Y \leq M$. First notice that $(S^{-1}y) \in Y$ but since

$$(S^{-1}y)_x = S^{-1}y - y \tag{6.8}$$

it turns out that $S^{-1}y$ has actually a C^1 representative (this is because H^1 is embedded in C^0). In order to check condition (1) then, we need just to show that $\sup_{x \in [0,1]} |(S^{-1}y)_x|$ is bounded by a constant depending only on M (see [18], p. 38). On the other hand, by (6.8)

$$\sup_{x \in [0,1]} |(S^{-1}y)_x| \leq \|S^{-1}y\|_{L^\infty} + \|y\|_{L^\infty} \leq c\|S^{-1}y\|_{H^1} + c\|y\|_{H^1},$$

where c is the constant entering in the Sobolev inequality corresponding to the inclusion $H^1 \hookrightarrow L^\infty$. Since S is an isometry we have $\|S^{-1}y\|_{H^1} = \|y\|_{L^2}$ and $\|y\|_{L^2} \leq \|y\|_{H^1}$ from which we deduce

$$\sup_{x \in [0,1]} |(S^{-1}y)_x| \leq 2cM. \tag{6.9}$$

Therefore, A is quasi-m-accretive on L^2 with ω bounded by cM .

To check assumption (2) we first show that $A(y)$ is a bounded operator from H^1 to L^2 for every $y \in H^1$. Indeed we have, for $w \in H^1$:

$$\|A(y)w\|_{L^2} = \|(S^{-1}y)w_x\|_{L^2} \leq \|S^{-1}y\|_{L^\infty} \|w_x\|_{L^2} \leq c\|S^{-1}y\|_{H^1} \|w\|_{H^1} =$$

$$= c\|y\|_{L^2}\|w\|_{H^1} \leq c\|y\|_{H^1}\|w\|_{H^1}.$$

To conclude the verification of (2), namely showing that (6.2) is fulfilled, we just observe that

$$\begin{aligned} \|(A(y) - A(z))w\|_{L^2} &= \|S^{-1}(y - z)w_x\|_{L^2} \leq \|S^{-1}(y - z)\|_{L^\infty}\|w_x\|_{L^2} \leq \\ &\leq c\|S^{-1}(y - z)\|_{H^1}\|w\|_{H^1} = c\|y - z\|_{L^2}\|w\|_{H^1}. \end{aligned}$$

Let us now verify the estimate (6.3). We notice that

$$SA(y)v - A(y)Sv = -(S^{-1}y)_x v_x,$$

then, for all $y, w \in H^1$, with $\|y\|_{H^1} \leq M$ we have

$$\|(SA(y) - A(y)S)S^{-1}w\|_{L^2} = \|-(S^{-1}y)_x(S^{-1}w)_x\|_{L^2},$$

which in virtue of (6.8) becomes

$$\|(S^{-1}y - y)(S^{-1}w - w)\|_{L^2}.$$

Therefore,

$$\begin{aligned} \|(SA(y) - A(y)S)S^{-1}w\|_{L^2} &= \|(S^{-1}y)(S^{-1}w) + yw - (S^{-1}w)y - (S^{-1}y)w\|_{L^2} \leq \\ &\leq \|(S^{-1}w)(S^{-1}y - y)\|_{L^2} + \|w(S^{-1}y - y)\|_{L^2} \leq (\|S^{-1}w\|_{L^2} + \|w\|_{L^2}) (\|S^{-1}y\|_{L^\infty} + \|y\|_{L^\infty}). \end{aligned}$$

Since $\|S^{-1}y\|_{L^\infty} \leq c\|S^{-1}y\|_{H^1} = c\|y\|_{L^2} \leq c\|y\|_{H^1}$, $\|y\|_{L^\infty} \leq c\|y\|_{H^1}$ and $\|S^{-1}w\|_{L^2} \leq \|S^{-1}y\|_{H^1} = \|y\|_{L^2}$, we get

$$\|(SA(y) - A(y)S)S^{-1}w\|_{L^2} \leq 4c\|y\|_{H^1}\|w\|_{L^2} \leq 4cM\|w\|_{L^2}.$$

In order to check the assumption (4), we first show $f(y) := -y(S^{-1}y)_x$ is bounded from $L := \{y \in H^1 : \|y\|_{H^1} \leq M\}$ to H^1 . We have

$$\|y(S^{-1}y)_x\|_{H^1}^2 = \|y(S^{-1}y)_x\|_{L^2}^2 + \|\partial_x(y(S^{-1}y)_x)\|_{L^2}^2,$$

and

$$\|y(S^{-1}y)_x\|_{L^2}^2 \leq \|(S^{-1}y)_x\|_{L^\infty}^2\|y\|_{L^2}^2 \leq 4c^2M^4,$$

in virtue of (6.9) and $\|y\|_{L^2} \leq \|y\|_{H^1} \leq M$. To control the other term, we observe that

$$\begin{aligned} \|\partial_x(y(S^{-1}y)_x)\|_{L^2} &\leq \|y_x(S^{-1}y)_x\|_{L^2} + \|y(S^{-1}y)_{xx}\|_{L^2} \leq \\ &\leq \|(S^{-1}y)_x\|_{L^\infty}\|y_x\|_{L^2} + \|y(S^{-1}y - y - y_x)\|_{L^2} \leq 2cM^2 + \|y\|_{L^\infty}(\|S^{-1}y\|_{L^2} + \|y\|_{L^2} + \|y_x\|_{L^2}) \leq \\ &\leq 2cM^2 + cM(3M) = 5cM^2, \end{aligned}$$

where we have used once again (6.9) and $\|y_x\|_{L^2} \leq \|y\|_{H^1} \leq M$. Combining these estimates we obtain

$$\|y(S^{-1}y)_x\|_{H^1}^2 \leq 4c^2M^4 + 25c^2M^4 = 29c^2M^4.$$

Let us prove the first Lipschitz inequality (6.4) is fulfilled. We have

$$\|f(y) - f(z)\|_{L^2} \leq \|yS^{-1}y - zS^{-1}z\|_{L^2} + \|y^2 - z^2\|_{L^2}.$$

For the first term $\|yS^{-1}y - zS^{-1}z\|_{L^2}$, adding and subtracting $zS^{-1}y$ we obtain

$$\|yS^{-1}y - zS^{-1}z\|_{L^2} \leq \|S^{-1}y\|_{L^\infty}\|y - z\|_{L^2} + \|S^{-1}(y - z)\|_{L^\infty}\|z\|_{L^2} \leq \quad (6.10)$$

$$\leq \|S^{-1}y\|_{H^1}\|y - z\|_{L^2} + \|S^{-1}(y - z)\|_{H^1}\|z\|_{L^2} = (\|y\|_{L^2} + \|z\|_{L^2})\|y - z\|_{L^2} \quad (6.11)$$

$$\leq \mu\|y - z\|_{L^2}, \quad (6.12)$$

where $\mu = \max\{\|y\|_{L^2}, \|z\|_{L^2}\}$. Notice now that we can not bound the term $\|y^2 - z^2\|_{L^2}$ with a Lipschitz constant depending only on the norms $\|y\|_{L^2}$ and $\|z\|_{L^2}$, but we will have to use the L^∞ -norms that are bigger. This is still sufficient, as one can see in the second inequality at page 40 of [18]. Then it is immediate to get

$$\|y^2 - z^2\|_{L^2} \leq (\|y\|_{L^\infty} + \|z\|_{L^\infty})\|y - z\|_{L^2}. \quad (6.13)$$

Combining this with the inequality controlling $\|yS^{-1}y - zS^{-1}z\|_{L^2}$, (6.4) is checked.

Finally we prove the last Lipschitz inequality (6.5). We have for every $y, z \in H^1$

$$\|y(S^{-1}y)_x - z(S^{-1}z)_x\|_{H^1} \leq \|y(S^{-1}y) - z(S^{-1}z)\|_{H^1} + \|y^2 - z^2\|_{H^1}, \quad (6.14)$$

using (6.8). We first control the first term on the right:

$$\begin{aligned} \|y(S^{-1}y) - z(S^{-1}z)\|_{H^1} &\leq \|y(S^{-1}y) - z(S^{-1}z)\|_{L^2} + \|\partial_x(y(S^{-1}y) - z(S^{-1}z))\|_{L^2} \leq \\ &\leq \|y(S^{-1}y) - z(S^{-1}z)\|_{L^2} + \|y_x(S^{-1}y) - z_x(S^{-1}z)\|_{L^2} + \|y(S^{-1}y)_x - z(S^{-1}z)_x\|_{L^2} \leq \\ &\leq (\|y\|_{L^2} + \|z\|_{L^2})\|y - z\|_{L^2} + \|y_x(S^{-1}y) - z_x(S^{-1}z)\|_{L^2} + \\ &\quad + (\|y\|_{L^2} + \|z\|_{L^2})\|y - z\|_{L^2} + (\|y\|_{L^\infty} + \|z\|_{L^\infty})\|y - z\|_{L^2}, \end{aligned}$$

using the inequalities (6.10), (6.11) and (6.13). Therefore we obtain for the first term on the right hand side of (6.14)

$$\|y(S^{-1}y) - z(S^{-1}z)\|_{H^1} \leq 3(\|y\|_{L^\infty} + \|z\|_{L^\infty})\|y - z\|_{H^1} + \|y_x(S^{-1}y) - z_x(S^{-1}z)\|_{L^2}.$$

On the other hand, adding and subtracting $y_x(S^{-1}z)$ inside $\|y_x(S^{-1}y) - z_x(S^{-1}z)\|_{L^2}$ we obtain

$$\|y_x(S^{-1}y) - z_x(S^{-1}z)\|_{L^2} \leq \|y_x(S^{-1}(y - z))\|_{L^2} + \|(S^{-1}z)(y_x - z_x)\|_{L^2} \leq$$

$$\begin{aligned}
&\leq \|S^{-1}(y-z)\|_{L^\infty} \|y_x\|_{L^2} + \|S^{-1}z\|_{L^\infty} \|y_x - z_x\|_{L^2} \leq \\
&\leq c\|S^{-1}(y-z)\|_{H^1} \|y\|_{H^1} + c\|S^{-1}z\|_{H^1} \|y-z\|_{H^1} \leq c(\|y\|_{H^1} + \|z\|_{H^1}) \|y-z\|_{H^1}.
\end{aligned}$$

This proves that the first term on the right hand side of (6.14) is bounded by $\mu\|y-z\|_{H^1}$ where μ depends only on $\max\{\|y\|_{H^1}, \|z\|_{H^1}\}$. Finally to control the second term on the right hand side of (6.14) observe that

$$\begin{aligned}
&\|y^2 - z^2\|_{H^1} \leq \|(y+z)(y-z)\|_{L^2} + \|\partial_x((y-z)(y+z))\|_{L^2} \leq \\
&\leq \|(y+z)(y-z)\|_{L^2} + \|(y-z)_x(y+z)\|_{L^2} + \|(y-z)(y+z)_x\|_{L^2} \leq \\
&\leq \|y+z\|_{L^\infty} \|y-z\|_{L^2} + \|(y-z)_x\|_{L^2} \|y+z\|_{L^\infty} + \|y-z\|_{L^\infty} \|(y+z)_x\|_{L^2} \leq \\
&\leq c(\|y\|_{H^1} + \|z\|_{H^1}) \|y-z\|_{H^1} + c\|y-z\|_{H^1} (\|y\|_{H^1} + \|z\|_{H^1}) + c\|y-z\|_{H^1} (\|y\|_{H^1} + \|z\|_{H^1}) \leq \\
&\leq 3c(\|y\|_{H^1} + \|z\|_{H^1}) \|y-z\|_{H^1}.
\end{aligned}$$

Therefore we can conclude that

$$\|y(S^{-1}y)_x - z(S^{-1}z)_x\|_{H^1} \leq \mu_3 \|y-z\|_{H^1},$$

where μ_3 depends only on $\max\{\|y\|_{H^1}, \|z\|_{H^1}\}$ so assumption (4) is satisfied and the Theorem is proved. $\blacksquare$

Remark 6.3 *In general the time T in the Theorem 6.2 depends on the norm $\|\phi\|_{H^1}$, hence for each choice of $M > 0$ such that $\|\phi\|_{H^1} \leq M$ we will have a corresponding T depending on M (see [18]). Moreover, since the operator $A(y)$ used in the above proof generates a C^0 -semigroup, and not just $-A(y)$ (see for instance [32], Theorem 12.26) we have local well-posedness also forward in time for the original equation (5.1).*

7 Numerical solutions

The present section is devoted to the numerical study of the initial value problem for the equation (5.1) in the case of periodic boundary conditions. Let us observe that introducing the auxiliary variable $P(x, t)$ such that

$$(1 - \partial_x)P = \frac{1}{2}v^2, \tag{7.1}$$

equation (5.1) can be equivalently written as follows

$$(1 - \partial_x)(v_t - vv_x - P_x) = 0. \tag{7.2}$$

Since the only smooth function f fulfilling periodic boundary conditions such that $(1 - \partial_x)f = 0$ is $f(x, t) \equiv 0$, the equation (7.2) is equivalent (on smooth solutions) to the system

$$\begin{aligned} v_t &= vv_x + P_x, \\ P_x &= P - \frac{1}{2}v^2. \end{aligned} \tag{7.3}$$

Numerical solutions are obtained by using a pseudo spectral method implemented in the `NDSolve Package` of `Mathematica 9` [34]. Although local existence of solutions is guaranteed by the Theorem 6.2, we observe that positive definite initial data seems to support global existence (in time) of solutions. Contrarily, finite-time break up occurs in other cases. In Figure 2 we compare the evolution at different time steps of the positive definite sinusoidal initial data

$$v_1(x) := v(x, 0) = \sin\left(\frac{\pi x}{12}\right) + 2 \tag{7.4}$$

$$v_2(x) := v(x, 0) = \sin\left(\frac{\pi x}{6}\right) + 2. \tag{7.5}$$

From the definition of $P(x, t)$ we have respectively

$$\begin{aligned} P_1(x) := P(x, 0) &= \frac{9}{4} - \frac{9 \cos\left(\frac{1}{6}\pi x\right)}{36 + \pi^2} + \frac{3\pi \sin\left(\pi x\right)}{2(36 + \pi^2)} + \frac{24\pi \cos\left(\frac{1}{12}\pi x\right)}{144 + \pi^2} + \frac{288 \sin\left(\frac{1}{12}\pi x\right)}{144 + \pi^2} \\ P_2(x) := P(x, 0) &= \frac{12\pi \cos\left(\frac{\pi x}{6}\right)}{\pi^2 + 36} - \frac{9 \cos\left(\frac{\pi x}{3}\right)}{4(\pi^2 + 9)} + \frac{72 \sin\left(\frac{\pi x}{6}\right)}{\pi^2 + 36} + \frac{3\pi \sin\left(\frac{\pi x}{3}\right)}{9(\pi^2 + 9)} + \frac{9}{4}. \end{aligned}$$

In both cases, after the initial steepening both solutions tend to vanish in time. It should be noted that the steepening is more pronounced for the datum of larger period (7.4) as expected, although the effect of dissipation prevents it to reach the gradient catastrophe. In the case of the Camassa-Holm equation it was proved that if the quantity $u(x, 0) - u_{xx}(x, 0)$ does not change sign, the periodic Cauchy problem is well-posed in H^2 [7]. It would be interesting to prove a similar results for the equation (5.1).

A completely different scenario takes place when considering initial data that change sign, as for instance an odd initial datum as

$$v_3(x) := v(x, 0) = \sin\left(\frac{\pi x}{12}\right).$$

The corresponding initial function P is

$$P_3(x) := P(x, 0) = \frac{3\pi \sin\left(\frac{\pi x}{6}\right)}{2(\pi^2 + 36)} - \frac{9 \cos\left(\frac{\pi x}{6}\right)}{\pi^2 + 36} + \frac{1}{4}.$$

As Figure 3 shows no damping occurs in this case and the solution seems to develop a gradient catastrophe in finite time at the inflection point $x = 0$. A similar phenomenon has been observed in the case of Camassa-Holm equation for which there




Figure 2: Time evolution for the initial datum $v(x, 0) = \sin\left(\frac{\pi x}{12}\right) + 2$ (solid line) and $v(x, 0) = \sin\left(\frac{\pi x}{6}\right) + 2$ (dashed line) with periodic boundary conditions.

exists a class of initial conditions $(u(x, 0) \in H^4$ with $u(x, 0)$ odd and $u(x, 0) \neq 0$) such that the solution blows up in finite time[6, 7].

Remark 7.1 *Let us observe that if G is the Green function for the operator $(1 - \partial_x)$ on the circle, then from the definition of the function P (7.1) we have $P = G \star \frac{1}{2}v^2$, where $\star$ denotes the convolution product. In full analogy with the Camassa-Holm equation, using this representation for P the equation (5.1) can be written as a conservation law with nonlocal flux*

$$v_t = \partial_x \left(\frac{1}{2}v^2 + G \star \frac{1}{2}v^2 \right).$$

8 Quasitriviality, transport equations and deformed hodograph formula

A Miura transformation is defined as a change of variables of the form

$$u(v, v_x, \dots) = \sum_k \epsilon^k F_k(v, v_x, v_{xx}, \dots), \quad \deg F_k = k \quad (8.1)$$

where coefficients F_k are differential polynomials in the derivatives $v_x, v_{xx}, \dots$. A natural generalisation of (8.1), referred to as *quasi-Miura* transformation, is obtained just relaxing the assumption on the polynomial form of F_k .




Figure 3: Steepening of the solution for the initial datum $v(x, 0) = \sin\left(\frac{\pi x}{12}\right)$.

According to the results of [23], any evolutionary PDEs of the form

$$v_t = vv_x + \sum_k \epsilon^k P_k(v, v_x, v_{xx}, \dots), \quad \deg P_k = k$$

can be reduced by a *quasi-Miura* transformation to the Hopf equation (3.2). Moreover any evolutionary symmetry of the equation

$$v_\tau = c(v)v_x + \sum_k \epsilon^k Q_k(v, v_x, v_{xx}, \dots)$$

is reduced by the same transformation to its dispersionless limit

$$u_\tau = c(u)u_x.$$

In particular, given an integrable hierarchy of evolutionary PDEs, there exists an invertible quasi-Miura transformation that brings any equation of the hierarchy to the corresponding equation of the Hopf hierarchy.

The method for constructing recursively the terms of a quasi-Miura transformation at any order in ϵ has been illustrated by S. Liu and Y. Zhang in [23] and it is based on the construction of infinitesimal generators $X_1, X_2, \dots$ for the reducing transformation

$$v(u) = \exp \tilde{X}(u) = u + \tilde{X}(u) + \frac{1}{2} \tilde{X}(\tilde{X}(u)) + \quad (8.2)$$

where

$$\tilde{X} = X \frac{\partial}{\partial u} + X_x \frac{\partial}{\partial u_x} + \dots$$

and

$$X = \epsilon X_1 + \epsilon^2 X_2 + \dots \quad .$$

In the following, we propose an alternative but equivalent version of the Liu-Zhang approach that is based on the direct solution of transport equations for the class of equations of the form

$$v_t = \partial_x (v^2 + \epsilon a(v)v_x + \epsilon^2 b_1(v)v_{xx} + \epsilon^3 c_1(v)v_{xxx} + \dots) . \quad (8.3)$$

Looking for asymptotic solutions in power series of ϵ of the form

$$v = u + \epsilon v^1 + \epsilon^2 v^2 + \dots \quad (8.4)$$

where $u = v^0$, the equation (8.3) splits into a quasilinear equation for u_0 plus a set of transport equations

$$\begin{aligned} Lu &= 0 \\ L^* v^1 &= \partial_x (a(u)u_x) \\ L^* v^2 &= \partial_x ((v^1 + \partial_x a(u))v^1 + b_1(u)u_{xx}) \\ &\dots \end{aligned} \quad (8.5)$$

where

$$L = \partial_t - 2u\partial_x \quad L^* = \partial_t - \partial_x 2u$$

are, respectively, the first order differential operator of the Hopf flow and its adjoint. Hence, the problem of constructing the reducing quasi-Miura transformation is reduced to the classical problem of constructing the expansion of the form (8.4) via the transport equations (8.5). Such solutions are referred to as formal solutions to the equation (8.3). Let us observe that once a solution u to the first equation in (8.5) is known, higher order corrections $v^1, v^2, \dots$ are obtained by solving a sequence of linear PDEs with coefficients depending on u . Hence, the asymptotic formal solution of the form (8.4) can be interpreted as a transformation that brings any solution to the Hopf equation to a solution of the deformed equation (8.3)[13]. It turns out that for the class of equations under consideration such a transformation is of quasi-Miura type. Given a solution to the Hopf equation $Lu = 0$ that is implicitly given in terms of hodograph equation:

$$x + 2ut + f(u) = 0, \quad (8.6)$$

the n -th transport equation can be written in the form

$$L^* v^n = F_n(u, u_x), \quad (8.7)$$

where we have observed that using the differential consequences of the equation (8.6)

$$u_{xx} = f''(u)u_x^3, \quad u_{xxx} = f'''(u)u_x^4 + 3(f'')^2u_x^5 \quad \text{etc.} \quad (8.8)$$

the r.h.s can be written as a suitable function of u and u_x only. The general integral of the equation (8.7) is readily obtained as a function of the variables u and u_x of the form

$$v^n = p_n(u, u_x)u_x + h_n(u, u_x) \quad (8.9)$$

where

$$p_n(u, u_x) = \int^{u_x} \frac{1}{2\varphi^3} F_n(u, \varphi) d\varphi$$

and $h_n(u, u_x)$ is the general solution to the homogenous linear equation $L^*h_n = 0$. Using the method of characteristics one can show that

$$h_n(u, u_x) = g_n(u)u_x$$

where $g_n(u)$ is an arbitrary function of its argument.

For the Burgers equation, that is obtained from (8.3) by choosing a constant central invariant $a(u) = 1$ the set of transport equation takes the a simple recursive form

$$\begin{aligned} L^*v^1 &= \partial_x(u_x) \\ L^*v^N &= \partial_x \left(\sum_{i=1}^{N-1} v^i v^{N-1-i} + v_x^{N-1} \right), \quad N = 2, 3, \dots \end{aligned}$$

with the notation $v^0 = u$.

Using the argument above one can prove that the solution to the n -th transport equation for the Burgers equation is written as a polynomial in u_x of the form

$$v^n = \sum_{j=1}^{2n-1} \alpha_{n,n+j} u_x^{n+j} + g_n(u)u_x \quad (8.10)$$

where coefficients α are determined by recursion as follows

$$\begin{aligned}
\alpha_{n,n+1} &= \frac{1}{2n} \alpha''_{n-1,n} \\
\alpha_{n,n+2} &= \frac{1}{2(n+1)} [\alpha''_{n-1,n+1} + \Lambda'_{n-1,2} + \Omega_{n-1,1}] \\
\alpha_{n,n+j} &= \frac{1}{2(n+j-1)} [\alpha''_{n-1,n+j-1} + \Lambda'_{n-1,j} + (n+j-1)f''(u)\Lambda_{n-1,j-1} + \Omega_{n-1,j-1} \\
&\quad + (n+j-3)(n+j-1)(f''(u))^2\alpha_{n-1,n+j-3}], \quad j = 3, \dots, 2n-3 \\
\alpha_{n,3n-2} &= \frac{1}{2(3n-3)} [\Lambda'_{n-1,2n-2} + (3n-3)f''(u)\Lambda_{n-1,2n-3} + \Omega_{n-1,2n-3} \\
&\quad + (3n-5)(3n-3)(f''(u))^2\alpha_{n-1,3n-5}] \\
\alpha_{n,3n-1} &= \frac{1}{2(3n-2)} [(3n-2)f''(u)\Lambda_{n-1,2n-2} + 3n(f''(u))^2\alpha_{n-1,3n-4}]
\end{aligned}$$

where

$$\begin{aligned}
\alpha_{1,2} &= \frac{1}{2} f''(u) \\
\Lambda_{n-1,k} &= \sum_{i=1}^{n-1} \sum_{j=1}^{2i-1} \alpha_{i,i+j} \alpha_{n-i,n+k-i-j}, \\
\Omega_{n-1,j} &= (2n+2j-1)f''(u)\alpha'_{n-1,n+j-1} + (n+j-1)f'''(u)\alpha_{n-1,n+j-1}.
\end{aligned}$$

Let us note that coefficients α 's depend only on the variable u through the function $f''(u)$ and its higher derivatives. The quasi-Miura transformation for Burgers' equation can be recovered from the formula (8.10) by choosing $g_n(u) = 0$ and eliminating the dependence on function $f''(u)$ and its higher derivatives via the triangular system of the form (8.8). It turns out that terms of quasi-Miura transformation at any order are rational functions of u_x, u_{xx} etc. with no explicit dependence on u . For instance, up to $O(\epsilon^4)$, the quasi-Miura transformation for Burgers' equation takes the form

$$v = u + \epsilon \frac{u_{xx}}{2u_x} + \epsilon^2 \left(\frac{u_{xxx}}{8u_x^2} - \frac{u_{xx}^2}{6u_x^3} \right)_x + \epsilon^3 \left(\frac{u_{5x}}{48u_x^3} - \frac{u_{xx}u_{4x}}{6u_x^4} - \frac{u_{xxx}^2}{8u_x^4} + \frac{3u_{xx}^2u_{xxx}}{4u_x^5} \right)_x + O(\epsilon^4). \quad (8.11)$$

A direct comparison with the Liu-Zhang approach to the solution of transport equations [23] is readily made by observing that the functions $f''(u), f'''(u), \dots$ play the role of parameters in the quadrature formula (8.9) as well as the functions $x_{uu} = f''_u, x_{uuu} = f'''_u, \dots$ do in Liu-Zhang's framework.

Theorem 8.1 *Given any solution to the Hopf equation*

$$u_t = 2uu_x,$$

via the hodograph formula (8.6), the formal solution of the Burgers equation of the form (8.11) satisfies the deformed hodograph equation

$$x + 2ut + \omega_f = 0 \quad (8.12)$$

where ω_f is the deformed 1-form corresponding to $a(u) = 1$.

Proof. Let us consider the vector field

$$P\alpha_f = \partial_x(x + 2ut + f) = 1 + 2u_x t + \partial_x f. \quad (8.13)$$

It can be viewed as a linear t -dependent combination of the vector fields

$$\frac{\partial}{\partial u}, u_x \frac{\partial}{\partial u}, (\partial_x f) \frac{\partial}{\partial u}.$$

Applying to such vector fields the inverse of the quasi-Miura transformation (8.11)

$$u = v + \frac{1}{2}\epsilon \frac{v_{xx}}{v_x} + \epsilon^2 \left(\frac{1}{8} \frac{v_{(4)}}{v_x^2} - \frac{7}{12} \frac{v_{xxx}v_{xx}}{v_x^3} + \frac{1}{2} \frac{v_{xx}^3}{v_x^4} \right) + \mathcal{O}(\epsilon^3) \quad (8.14)$$

and taking into account the transformation law for vector fields

$$X(u) \rightarrow \tilde{X}(v) = \left(\frac{\partial v}{\partial u} + \frac{\partial v}{\partial u_x} \partial_x + \frac{\partial v}{\partial u_{xx}} \partial_x^2 + \dots \right) X(u)|_{u=u(v, v_x, \dots)}$$

we can immediately prove that

- $\frac{\partial}{\partial u} \rightarrow \frac{\partial}{\partial v}$ as a consequence of the fact that $\frac{\partial v}{\partial u} = 1$.
- $u_x \frac{\partial}{\partial u} \rightarrow v_x \frac{\partial}{\partial v}$.
- $(\partial_x f) \frac{\partial}{\partial u} \rightarrow (\partial_x \omega_f) \frac{\partial}{\partial v}$. In the general case we only know that such a deformation exists up to the order ϵ^5 , but in the case of Burgers it is defined for an arbitrary (analytic) function f .

Combining the above results we obtain

$$(1 + 2u_x t + \partial_x f) \frac{\partial}{\partial u} \rightarrow (1 + 2v_x t + \partial_x \omega_f) \frac{\partial}{\partial v}$$

This means that, given any hodograph solution $u(x, t)$ the series (8.14) satisfies the equation

$$1 + 2v_x t + \partial_x \omega_f = 0. \quad (8.15)$$

Integrating with respect to x we obtain

$$x + 2v t + \omega_f = c. \quad (8.16)$$

where c is a constant. Taking the limit $\epsilon \rightarrow 0$ it is immediate to check that the constant c must vanish. ■

Let us now consider the general case where the function a is not constant and suppose, as we have conjectured, that there exists an integrable hierarchy for any choice of the central invariant $a(u)$. As a consequence of the results of S. Liu and Y. Zhang there should exist also in this case a reducing quasi-Miura transformation. One can easily check that such a transformation depends in general on v at any order in the deformation parameter. This immediately implies that the vector field $\frac{\partial}{\partial u}$ is no longer invariant and an additional contribution must be taken into account. If this additional term is a total x -derivative, one obtains a correction to the deformed hodograph formula (8.12). Indeed if

$$\frac{\partial v}{\partial u} = 1 + \partial_x (F(u))$$

then $\frac{\partial}{\partial u} \rightarrow [1 + \partial_x (F(u(v)))] \frac{\partial}{\partial v}$ and therefore

$$(1 + 2u_x t + \partial_x f) \frac{\partial}{\partial u} \rightarrow (1 + 2v_x t + \partial_x (\omega_f + F(u(v)))) \frac{\partial}{\partial v}.$$

Proceeding exactly as in the above theorem one obtains the deformed hodograph formula

$$x + 2v t + \omega_f + F = 0. \quad (8.17)$$

For example, in the case $a(u) = u$ the reducing transformation is

$$v = u + \frac{1}{2}\epsilon \left(\frac{uu_{xx}}{u_x} + u_x \ln u_x \right) + \frac{1}{4}\epsilon^2 \left(\frac{1}{2} \frac{u^2 u_{(4)}}{u_x^2} + 3 \frac{uu_{xxx}}{u_x} - \frac{7}{3} \frac{u^2 u_{xxx} u_{xx}}{u_x^3} - \frac{7}{3} \frac{uu_{xx}^2}{u_x^2} + 2 \frac{u^2 u_{xx}^3}{u_x^4} + \frac{1}{2} u_{xx} (\ln u_x)^2 + 2u_{xx} \ln u_x + 2u_{xx} + \frac{uu_{xxx} \ln u_x}{u_x} - \frac{uu_{xx}^2 \ln u_x}{u_x^2} \right) + \mathcal{O}(\epsilon^3)$$

whose inverse is

$$u = v - \frac{1}{2}\epsilon \left(\frac{vv_{xx}}{v_x} + v_x \ln v_x \right) - \epsilon^2 \left(-\frac{1}{8} \frac{v^2 v_{(4)}}{v_x^2} - \frac{1}{4} \frac{vv_{xxx}}{v_x} + \frac{5}{12} \frac{v^2 v_{xxx} v_{xx}}{v_x^3} - \frac{1}{12} \frac{vv_{xx}^2}{v_x^2} - \frac{1}{4} \frac{v^2 v_{xx}^3}{v_x^4} - \frac{1}{8} v_{xx} (\ln v_x)^2 - \frac{1}{2} v_{xx} \ln v_x - \frac{1}{4} \frac{vv_{xxx} \ln v_x}{v_x} + \frac{1}{4} \frac{vv_{xx}^2 \ln v_x}{v_x^2} \right) + \mathcal{O}(\epsilon^3).$$

After some computations one gets

$$\begin{aligned} \frac{\partial v}{\partial u} &= 1 + \partial_x \left[\frac{1}{2}\epsilon \ln u_x + \epsilon^2 \left(\frac{1}{4} \frac{uu_{xxx}}{u_x^2} + \frac{1}{4} \frac{u_{xx} \ln u_x}{u_x} + \frac{1}{2} \frac{u_{xx}}{u_x} - \frac{1}{3} \frac{uu_{xx}^2}{u_x^3} \right) \right] + \mathcal{O}(\epsilon^3) = \\ &= 1 + \partial_x \left[\frac{1}{2}\epsilon \ln v_x - \epsilon^2 \left(\frac{1}{12} \frac{vv_{xx}^2}{v_x^3} \right) \right] + \mathcal{O}(\epsilon^3) \end{aligned}$$

and therefore the correction to the deformed hodograph formula up to the second order is given by

$$F(v) = \frac{1}{2}\epsilon \ln v_x - \epsilon^2 \left(\frac{1}{12} \frac{vv_{xx}^2}{v_x^3} \right).$$

The asymptotic approach based on the construction of formal solutions via quasi-Miura transformations or equivalently via transport equations is generally expected to provide an accurate local asymptotic description of solutions to equations of the form (8.3) for sufficiently regular initial data and sufficiently small times, i.e. before the generic solution to the Hopf equation $Lu = 0$ develops a gradient catastrophe. The rigorous justification of this method requires a dedicated analysis that goes beyond the scope of the present work. However, for some equations of the form (8.3) rigorous results are already available (see e.g. [25, 26]). As it was pointed out in [26] the fact that the quasi-Miura transformation (8.1) does not preserve the initial datum makes difficult to perform a direct comparison between the perturbed and the unperturbed solution to a specific initial value problem. This problem can be readily fixed using the fact that terms at any order in ϵ of a quasi-Miura transformation are particular solutions to transport equations (8.5) whose general integral is defined up to the kernel of the adjoint operator L^* . Hence, given a solution to the Hopf equation $Lu = 0$ with initial datum $u(x, 0) = u_0(x)$, the required perturbed solution $v = u + \epsilon v^1 + \epsilon^2 v^2 + \dots$ that satisfies the same initial datum has to be such that $v^1(x, 0) = 0$, $v^2(x, 0) = 0$ etc. Such solutions are obtained by choosing the function $h_n(u, u_x) = g_n(u)u_x$ in (8.9) in such a way that

$$g_n(u_0) = -p_n(u_0, u_{0x}) = -p_n\left(u_0, -\frac{1}{f'(u_0)}\right)$$

where we have used the formula

$$u_{0x} = -\left.\frac{1}{2t + f'(u)}\right|_{t=0} = -\frac{1}{f'(u_0)}.$$

In the KdV case the existence of a transformation reducing KdV equation to Hopf equation and preserving the initial data was proved in [25]. The above arguments show that, in general, the existence of this transformation relies on the freedom in the choice of solutions of transport equations (8.7).

9 Dubrovin's Universality

Let us consider the Hopf equation, that is the conservation law associated with the undeformed 1-form $\omega_{u^2} = u^2$

$$u_t = 2uu_x, \tag{9.1}$$

and its solution given in terms of the formula

$$x + 2ut - f(u) = 0 \tag{9.2}$$

where the function $f(u)$ is an arbitrary function that parametrizes the family of commuting flows associated with the unperturbed 1-form $\omega_f = f(u)$

$$u_\tau = f'(u)u_x.$$

The deformation procedure proposed leads to the pair of involutive 1-forms

$$\begin{aligned}\omega_{u^2}^{def} &= u^2 + \epsilon a u_x + \epsilon^2 a a' u_{xx} + O(\epsilon^3) \\ \omega_f^{def} &= f(u) + \frac{\epsilon}{2} a f'' u_x + \epsilon^2 \left[\left(\frac{1}{2} a a' f''' + \frac{1}{6} a^2 f''' \right) u_{xx} + \left(\frac{1}{2} a a' f''' + \frac{1}{8} a^2 f^{(4)} \right) u_x^2 \right] + O(\epsilon^3)\end{aligned}$$

where $a = a(u)$. Let us now proceed with the study of the critical behaviour of Burgers equation in full analogy with the case of dispersive Hamiltonian equations considered by Dubrovin in [12]. Introducing the deformed hodograph equation

$$x + 2ut - \omega_f^{def} = 0 \quad (9.3)$$

where ω_f^{def} has been specified for a constant central invariant $a(u) = a_0$, we shall perform a multiscale analysis about the generic point of gradient catastrophe (x_0, t_0, u_0) such that

$$x_0 + 2u_0 t_0 - f(u_0) = 0 \quad 2t_0 - f'(u_0) = 0 \quad f''(u_0) = 0 \quad f'''(u_0) > 0. \quad (9.4)$$

Introducing displacement variables $(\bar{x}, \bar{t}, \bar{u})$ as follows

$$x = x_0 + \lambda^\alpha \bar{x} \quad t = t_0 + \lambda^\beta \bar{t} \quad u = u_0 + \lambda \bar{u} \quad (9.5)$$

where $\lambda = \epsilon^q$ is a small parameter, let us now expand in Taylor series the l.h.s of equation (9.3). Using the conditions (9.4) one obtains

$$\begin{aligned}\epsilon^{q\sigma} \tilde{x} + 2\epsilon^{q(\beta+1)} \bar{u} \bar{t} - \frac{\epsilon^{3q}}{6} f_0''' \bar{u}^3 - \frac{\epsilon^{1+q(2-\sigma)}}{2} a_0 f_0''' \bar{u} \bar{u}_{\tilde{x}} - \frac{\epsilon^{2+q(1-2\sigma)}}{6} a_0^2 f_0''' \bar{u}_{\tilde{x}\tilde{x}} \\ = O(\epsilon^{4q}) + O(\epsilon^{1+q(2-\sigma)}) + O(\epsilon^{2+q(2-2\sigma)})\end{aligned} \quad (9.6)$$

where the variable $\tilde{x}$ is defined as

$$\lambda^\sigma \tilde{x} = \lambda^\alpha \bar{x} + 2u_0 \lambda^\beta \bar{t} = x - x_0 + 2u_0(t - t_0),$$

with the notation $f_0 = f(u_0)$ etc. The request that all terms into the l.h.s of (9.6) contribute to the same order in ϵ gives

$$\sigma = 3 \quad \beta = 2 \quad q = \frac{1}{4}.$$

Hence, we we end up with the expression

$$\tilde{x} + 2\bar{u}\bar{t} - \frac{1}{6} f_0''' \bar{u}^3 - \frac{1}{2} a_0 f_0''' \bar{u} \bar{u}_{\tilde{x}} - \frac{1}{6} a_0^2 f_0''' \bar{u}_{\tilde{x}\tilde{x}} = O(\epsilon^{1/4}). \quad (9.7)$$

Then, the solution to the deformed hodograph equation in the vicinity of the critical point satisfies the second order ODE obtained by taking the limit $\epsilon \rightarrow 0$ in (9.7). The above equation can conveniently be written in non-dimensional form via the following rescaling of the dependent and independent variables

$$\tilde{x} = s_1 X \quad \tilde{t} = s_2 T \quad \bar{u} = s_3 U$$

where

$$s_1 = \left(\frac{1}{6} a_0^3 f_0''' \right)^{1/4} \quad s_2 = \frac{1}{2} \frac{s_1^2}{a_0} = \left(\frac{1}{24} a_0 f_0''' \right)^{1/2} \quad s_3 = \frac{a_0}{s_1} = \left(\frac{6a_0}{f_0'''} \right)^{1/4}$$

and $U = U(X, T)$ satisfies the equation

$$U_{XX} + 3UU_X + U^3 - UT = X. \quad (9.8)$$

In terms of the original variables, the critical behaviour near the point of gradient catastrophe given a solution to the equation (9.8) is

$$u = u_0 + \left(\frac{6a_0}{f_0'''} \right)^{1/4} \epsilon^{1/4} U \left[\left(\frac{6}{a_0^3 f_0'''} \right)^{1/4} \frac{x - x_0 + 2u_0(t - t_0)}{\epsilon^{3/4}}, \left(\frac{24}{a_0 f_0'''} \right)^{1/2} \frac{t - t_0}{\epsilon^{1/2}} \right]. \quad (9.9)$$

We also observe that the equation (9.8) can be transformed via a Cole-Hopf transformation of the form $U = \partial_X \log w$ into the linear ODE

$$w_{XXX} - Tw_X = Xw. \quad (9.10)$$

Equation (9.10) has the following general solution

$$w(X, T) = c_1 {}_0F_2([1/2, 3/4], \frac{1}{64} (T + X)^4) + c_2 {}_0F_2([3/4, 5/4], \frac{1}{64} (T + X)^4) (T + X) \\ + c_3 {}_0F_2([5/4, 3/2], \frac{1}{64} (T + X)^4) (T + X)^2,$$

where c_1, c_2, c_3 are arbitrary constant of integration and ${}_0F_2$ is a generalized hypergeometric function defined by

$${}_0F_2([\alpha, \beta], z) := \sum_{n=0}^{+\infty} \frac{1}{(\alpha)_n (\beta)_n} \frac{z^n}{n!}, \quad (9.11)$$

where

$$(\alpha)_n := (\alpha)(\alpha + 1)(\alpha + 2) \dots (\alpha + n - 1), \quad (\alpha)_0 := 1$$

and analogously for $(\beta)_n$. The function ${}_0F_2([\alpha, \beta], z)$ defines an entire function whenever $\alpha > 0$ and $\beta > 0$ (it is clear from the expansion (9.11) that is dominated by the series of the exponential function up to a constant).

According with the conjecture formulated in [9] that generalises a result by A. Il'in [21], the critical behaviour (9.9) is provided by the particular solution to the equation (9.8)

$$U = \partial_X \log P(X, T) = \frac{P_X(X, T)}{P(X, T)} \quad (9.12)$$

where $P(X, T)$ is the Pearcey integral

$$P(X, T) = \int_{-\infty}^{\infty} e^{-(4z^4 - 2Tz^2 + 2Xz)} dz.$$

By a direct calculation using the identity

$$\int_{-\infty}^{\infty} (-16z^3 + 4Tz - 2X) e^{-(4z^4 - 2Tz^2 + 2Xz)} dz = 0$$

one can directly verify that the function $P(X, T)$ satisfies the linear ODE (9.10).

Acknowledgements

The authors wish to thank Roberto Camassa, Boris Dubrovin and Gregorio Falqui for useful discussions and references. The research of P.L. is partially supported by the Italian MIUR Research Project *Teorie geometriche e analitiche dei sistemi Hamiltoniani in dimensioni finite e infinite* and by GNFM Programme *Progetto Giovani 2012*. The research of A.M. is partially supported by the ERC grant *FroM-PDE*. A.M. is also grateful to B. Dubrovin and T. Grava for the kind hospitality at SISSA in Trieste.

References

- [1] A. Arsie and P. Lorenzoni, *Poisson bracket on 1-forms and evolutionary PDEs*, Journal of Physics A, Mathematical and Theoretical, (2012).
- [2] V.A. Baikov, R.K. Gazizov and N.Kh. Ibragimov, Approximate symmetries and formal linearization, J. Appl. Mech. Tech. Phys. 30, no. 2 (1989) 204-212.
- [3] H. Brezis, *Functional Analysis, Sobolev Spaces and Partial Differential Equations*, Springer-Verlag, (2011).
- [4] F. Calogero, *Why are certain nonlinear PDEs both widely applicable and integrable*, in “What is integrability”, V.E. Zakharov (Ed.), Springer-Verlag, Berlin-Heidelberg (1991), 1–62,.
- [5] M. Chen, S.Q. Liu, Y. Zhang, *A two-component generalization of the Camassa-Holm equation and its solutions*, Lett. Math. Phys. **75**(1), 1-15 (2006).

- [6] R. Camassa and D. Holm *An integrable shallow water wave equation with peaked solitons*, Physical Review Letters, **71**, no. 11, (1993).
- [7] A. Constantin, *On the Cauchy problem for the periodic Camassa-Holm equation*, J. of Diff. Equations, **141**, 218-235 (1997).
- [8] L. Degiovanni, F. Magri, V. Sciacca *On Deformation of Poisson Manifolds of Hydrodynamic Type* Commun. Math. Phys. **253**, 124 (2005).
- [9] B. Dubrovin, M. Elaeva, *On critical behaviour in nonlinear evolutionary PDEs with small viscosity*, Russian Journal of Mathematical Physics, vol. 19, p. 13-22.
- [10] B.A. Dubrovin, S.P. Novikov, *On Hamiltonian brackets of hydrodynamic type*, Soviet Math. Dokl. **279:2** (1984) 294–297.
- [11] B. Dubrovin, *Flat pencils of metrics and Frobenius manifolds*, Integrable systems and algebraic geometry (Kobe/Kyoto), 1997), 47-72, World Sci. Publishing, River Edge, NJ (1998).
- [12] B. Dubrovin, *Hamiltonian perturbations of hyperbolic systems of conservation laws II*, Comm. Math. Phys. Volume **267**, Number 1, 117-139, (2006).
- [13] B. Dubrovin, Y. Zhang, *Normal forms of integrable PDEs, Frobenius manifolds and Gromov-Witten invariants*, math.DG/0108160.
- [14] B. Dubrovin, S.-Q.Liu, Y.Zhang, *On hamiltonian perturbations of hyperbolic systems of conservation laws, I: quasitriviality of bihamiltonian perturbations*, Comm. Pure and Appl. Math. **59**, 559–615 (2006);
- [15] J. Escher, O. Lechtenfeld, Z. Yin, *Well-Posedness and blow-up phenomena for the 2-component Camassa-Holm equation*, Discrete Contin. Dyn. Syst. **19**(3), 493-513 (2007).
- [16] G. Falqui, *On a Camassa-Holm type equation with two dependent variables*, J. Phys. A: Math. Gen., **39**, 327–342 (2006).
- [17] A. S. Fokas, *A symmetry approach to exactly solvable evolution equations*, J. Math. Phys., **21**(6), 1318–1325 (1980).
- [18] T. Kato, *Quasi-linear equations of evolution, with applications to partial differential equations*, in “Spectral Theory and Differential Equations”, Springer Lecture Notes in Mathematics, Vol. 448, pp 25–70, Springer-Verlag, Berlin/New York, 1975.

- [19] E. Getzler, *A Darboux theorem for Hamiltonian operators in the formal calculus of variations*, Duke Math. J. **111** (2002) 535–560.
- [20] I.M. Gel'fand and I.A. Dorfman, *Hamiltonian operator and the classical Yang-Baxter equation*, Funct. Anal. Appl, 16 (1982), 241–248.
- [21] A.M. Il'in, *Matching of Asymptotic Expansions of Solutions of Boundary Value Problems*, AMS Translations of Mathematical Monographs, Vol. 102, (1992).
- [22] S-Q. Liu, Y. Zhang *Deformations of semisimple bihamiltonian structures of hydrodynamic type*, J. Geom. Phys. **54**(4), 427–453, (2005).
- [23] S-Q. Liu, Y. Zhang *On Quasitriviality and Integrability of a Class of Scalar Evolutionary PDEs* J. Geom. Phys. **57**, 101–119, (2006).
- [24] P. Lorenzoni, *Deformations of bi-Hamiltonian structures of hydrodynamic type*, J. Geom. Phys. **44** (2002) 331–375.
- [25] D. Masoero and A. Raimondo, *Semiclassical limit for generalized KdV equations before the gradient catastrophe*, arXiv:1107.0461v2 (2011).
- [26] D. Masoero, A. Raimondo, *A deformation of the method of characteristics and the Cauchy problem for Hamiltonian PDEs in the small dispersion limit*, arXiv:1211.2676 (2012).
- [27] A. V. Mikhailov, A. B. Shabat, and V. V. Sokolov, *The symmetry approach to classification of integrable equations*, In What is integrability? Springer Ser. Nonlinear Dynamics, 115184. Springer, Berlin (1991).
- [28] A. V. Mikhailov and V. S. Novikov, *Perturbative symmetry approach*, J. Phys. A, **35**(22), 4775–4790 (2002).
- [29] A. V. Mikhailov, A. B. Shabat, and R. I. Yamilov *A symmetric approach to the classification of nonlinear equations. Complete lists of integrable systems*, Uspekhi Mat. Nauk, **42**(4(256)) 3–53 (1987).
- [30] P. Olver, *Evolution equations possessing infinitely many symmetries*, J. of Mathematical Physics, **18**, 1212 (1977).
- [31] P. J. Olver, P. Rosenau, *Tri-Hamiltonian duality between solitons and solitary-wave solutions having compact support*, Phys. Rev. E, **53**(2), 1900–1906 (1996).
- [32] M. Renardy and R. C. Rogers, *An Introduction to Partial Differential Equations*, Texts in Applied Mathematics, **13**, Springer-Verlag (2003).

- [33] I.A.B.Strachan, *Deformations of the Monge/Riemann hierarchy and approximately integrable systems*, J. Math. Phys. **44** (2003) 251–262.
- [34] Wolfram Research, Inc. Mathematica 9 (Copyright 1988-2013).